

1)

The emergence Reinkeho edema significantly contributes to menopause. It is one of the most important factors. Our data suggest that it would not be so if they were women hormonally substituted.

The second most important moment for the recurrence of RE are manifestations of gastroesophageal reflux. It can be accompanied by reflux laryngofaryngeálním, but not necessarily. It seems

that its influence may be indirect, through vagal innervation (reflex). Uncover and reflux time it zaléčit proton pump is essential for proper treatment.

Diagnostic methods are different with regard to the patient's good to find the least burdensome method. Dynamic scintigraphy of the esophagus is in such examination.

Antibodies against HP in the stool indicate the need to treat helikobakteriózu regardless of the extent of gastrointestinal symptoms. Essential for initiation of drug treatment

finding Reinkeho edema as one of the manifestations of GERD. HP eradication can be achieved by the use of proton pump inhibitors. In refractory cases

the H + blocker or associated with typical gastroenterological problems, we recommend to treat the patient with antibiotics and chemotherapeutics

Smoking has a significant influence on the RE, especially if the patient stops smoking after surgery, relapse escapes with great hope. We do not at this point over

stress often inflected inappropriate use of cigarettes, but according to our findings, RE another of the many diseases that are triggered by smoking. Fight against

smoking support.

For edema is essential the correct use of voice. Phoniatic postoperative care decreases the risk of recurrent course significantly. Not all hospitals after

mikrolaryngeálním surgery are referred to the voice specialists. There is certainly a large therapeutic margin. Foniatra privileged position next to the surgeon's confirmed by our histological findings.

The possibility of recurrence outside the RE we have examined the factors involved and operatérova technique. We used classical instrumentation Kleinsasserovo. In

Currently, otolaryngology and penetrate new laser method (91), which has been used in the larynx. In these studies will be necessary to explore

therapeutic use ErCr: YSGG laser (Waterlase) with a wavelength of 2780 nm Nd: YVQ4 laser (Idas) the wavelength of 532 nm.

2)

The investigation and hoarseness Reinkeho edema is widely used traditional investigative techniques ENT. Besides indirektní laryngoscopy, magnifying

laryngoscopy and

stroboskopie, fibrolaryngoskopie and voice analysis is necessary to perform a one evaluated for the diagnosis of HP and GERD.

We have proved in the diagnosis of antibody tests of stool. Possible inhalation test. Both methods enable convenient and accurate diagnosis.

To pass GERD is sovereign by pH metry. This test is not rejected by the patient for their invasiveness. Dynamic scintigraphy of oesophageal pH can Metn substitute and is much gentler.

3)

The finding of synovial cells with similar cells in space Reinkeho convinces us of the correctness of the theory mikrotrauma in the pathogenesis of RE. In addition, our data confirm

significance and importance of proper phonation phoniatric spring. We care foniatra re ia after surgery is one of effective tools to combat recidivism RE.