

## ERRATA

- Str. 11** chybně uvedeno „Sarnander (1908)“ – správně: Sernander (1908)
- Str. 11** chybně uvedeno „Blytt-Sandarður model“ – správně: Blytt-Sernanderður model
- Str. 13** chybně uvedeno „Převzato z Walker et al. (1999)“ – správně: Převzato z Björck et al. (1998)
- Str. 17** chybně uvedeno „ve vozcích“ – správně: ve voscích
- Str. 17** chybně uvedeno „(Hou et al., 2008; Lindgreen & Lindgreen, 2004)“ – správně: (Hou et al., 2008; Huang et al., 2004)
- Str. 51** chybně uvedeno „(Obr. č. 20)“ – správně: (Obr. č. 19)
- Str. 54** chybně uvedeno „z Prášilského jezera“ – správně: z Plešného jezera
- Str. 57 a 70** chybně uvedeno „Jankovská, 2014“ – správně: Jankovská, 2006
- Str. 63 a 72** chybně uvedeno „Petr et al., 2015“ – správně: Vočadlova et al., 2015
- Str. 67** revize použité literatury:

## POUŽITÁ LITERATURA

- Alley, R. B. (2000). The Younger Dryas cold interval as viewed from central Greenland. *Quaternary Science Reviews*, 19, 213-236. [https://doi.org/10.1016/S0277-3791\(99\)00062-1](https://doi.org/10.1016/S0277-3791(99)00062-1)
- Ammann, B., Birks, H. J. B., Brooks, S. J., Eicher, U., Grafenstein, U. Von, Hofmann, W., Lemdahl, G., Schwander, J., Tobolski, K., & Wick, L. (2000). Quantification of biotic responses to rapid climatic changes around the Younger Dryas — a synthesis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 159, 313-347. [https://doi.org/10.1016/S0031-0182\(00\)00092-4](https://doi.org/10.1016/S0031-0182(00)00092-4)
- Andersson, G. (1896) Svenska växtvärldens historia. Stockholm: P.A. Norstedt & Soner. 136 pp.
- Arseneau, K. M. A., Driscoll, C. T., Brager, L. M., Ross, K. A., Cumming, B. F., & Weyhenmeyer, G. A. (2011). Recent evidence of biological recovery from acidification in the Adirondacks (New York, USA): a multiproxy paleolimnological investigation of Big Moose Lake. *Canadian Journal of Fisheries and Aquatic Sciences*, 68(4), 575–592. <https://doi.org/10.1139/f2011-003>
- Bastviken, D., Ejlertsson, J., & Tranvik, L. (2001). Similar bacterial growth on dissolved organic matter in anoxic and oxic lake water. *Aquatic Microbial Ecology*, 24(1), 41–49. <https://doi.org/10.3354/ame024041>
- Bennett, K. D. (1996). Determination of the number of zones in a biostratigraphical sequence. *New Phytologist*, 132(1), 155–170. <https://doi.org/10.1111/j.1469-8137.1996.tb04521.x>
- Birks, H. J. B., & Birks, H. H. (2008). Biological responses to rapid climate change at the Younger Dryas – Holocene transition at Kråkenes, western Norway. *The Holocene*, 18(1), 19–30. <https://doi.org/10.1177/0959683607085572>
- Björck, S., Walker, M. J. C., Cwynar, L. C., Johnsen, S., Knudsen, K. L., Lowe, J. J., & Wohlfarth, B. (1998). An event stratigraphy for the last termination in the North Atlantic region based on the Greenland ice-core record: a proposal by the INTIMATE group. *Journal of Quaternary Science*, 13(4), 283–292. [https://doi.org/10.1002/\(SICI\)1099-1417\(199807/08\)13:4<283::AID-JQS386>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1099-1417(199807/08)13:4<283::AID-JQS386>3.0.CO;2-A)
- Björck, S. (2013). Younger Dryas Oscillation, Global Evidence. In S. A. Elias, (Ed.): *Encyclopedia of Quaternary Science*, Volume 3, pp. 222–228. Elsevier B.V. Oxford.

- Błędzki, L. A., Szeroczyńska, K., & Puusepp, E. (2013). The late holocene appearance of European *Bosmina* (Eubosmina) thersites (Crustacea, Cladocera) in lakes surrounding the Baltic Sea. *Hydrobiologia*, 715(1), 77–86. <https://doi.org/10.1007/s10750-012-1376-6>
- Blytt, A. (1876). Essay on the Immigration of the Norwegian Flora During Alternating Rainy and Dry Periods. Christiania, 89 pp.
- Brauer, A., Endres, C., Gu, C., Litt, T., Stebich, M., & Negendank, F. W. (1999). High resolution sediment and vegetation responses to Younger Dryas climate change in varved lake sediments from Meerfelder Maar, Germany. *Quaternary Science Review*, 18, 321–329. [https://doi.org/10.1016/S0277-3791\(98\)00084-5](https://doi.org/10.1016/S0277-3791(98)00084-5)
- Broecker, S. (1998). Paleocean circulation during the Last Deglaciation: A bipolar seesaw? *Paleoceanography*, 13(2), 119–121.
- Brooks, S. J., & Birks, H. J. B. (2001). Chironomid-inferred air temperatures from Lateglacial and Holocene sites in north-west Europe: Progress and problems. *Quaternary Science Reviews*, 20(16), 1723–1741. [https://doi.org/10.1016/S0277-3791\(01\)00038-5](https://doi.org/10.1016/S0277-3791(01)00038-5)
- Brown, L. A. (1929). The Natural History of Cladocerans in Relation to Temperature-III. Preadaptation and Dispersal. *The American Naturalist*, 63(688), 443–454.
- Bubenková, A. (2018). Využití subfossilních perlooček v paleoekologických rekonstrukcích. Bakalářská práce, 40 str., Přírodovědecká fakulta, Univerzita Karlova, Praha
- Buizert, C., Adrian, B., Ahn, J., Albert, M., Alley, R. B., Baggenstos, D., Bauska, T. K., Bay, R. C., Bencivengo, B. B., Bentley, C. R., Brook, E. J., Chellman, N. J., Clow, G. D., Cole-Dai, J., Conway, H. et al. (2015). Precise inter-polar phasing of abrupt climate change during the last ice age. *Nature*, 520(7549), 661–665. <https://doi.org/10.1038/nature14401>
- Cao, M. (2016). Biomarker proxies as quantitative indicators of climate variability in the Montcortés lake. PhD Thesis, 183 pp., Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona
- Carlson, A. E. (2013). The Younger Dryas Climate Event. In: Elias S.A. (ed.) *Encyclopedia of Quaternary Science*, 3, 126–134. Elsevier B.V. <https://doi.org/10.1016/B978-0-444-53643-3.00029-7>
- Carvalho, M.A., Bengtson, P., Lana, C. C., Sá, N. de P., Santiago, G., & Giannerini, M. C. da S. (2019). Late Aptian (Early Cretaceous) dry–wet cycles and their effects on vegetation in the South Atlantic: Palynological evidence. *Cretaceous Research*, 100, 172–183. <https://doi.org/10.1016/j.cretres.2019.03.021>
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjörnsdóttir, A. E., Jouzel, J., & Bond, G. (1993). Evidence for general instability of past climate from a 250-kyr ice-core record. *Nature*, 364, 218–220.
- Dau, J. H. C. (1829). Allerunterthänigster Bericht an die Königliche Dänische Rentekammer über die Torfmoore Seelands nach einer im Herbste 1828 deshalb unternommenen Reise. *Kopenhagen und Leipzig: Gylendahl und Hinrichs*, 316 pp. <http://dx.doi.org/10.3931/e-rara-63273>
- Davison, W. (1993). Iron and manganese in lakes. *Earth Science Reviews*, 34, 119–163. [https://doi.org/10.1016/0012-8252\(93\)90029-7](https://doi.org/10.1016/0012-8252(93)90029-7)
- Dellwig, O., Leipe, T., März, C., Glockzin, M., Pollehne, F., Schnetger, B., Yakushev, E.V., Böttcher, M. E., & Brumsak, H. J. (2010). A new particulate Mn-Fe-P-shuttle at the redoxcline of anoxic basins. *Geochimica et Cosmochimica Acta*, 74(24), 7100–7115. <https://doi.org/10.1016/j.gca.2010.09.017>
- Delmas, M., Calvet, M., & Gunnell, Y. (2009). Variability of Quaternary glacial erosion rates - A global perspective with special reference to the Eastern Pyrenees. *Quaternary Science Reviews*, 28(5–6), 484–498. <https://doi.org/10.1016/j.quascirev.2008.11.006>
- Drzymulska, D. (2014). *Betula nana* L. (vaevakase) jääajajärgne levik Kirde-Poolas. *Estonian Journal of Earth Sciences*, 63(2), 76–87. <https://doi.org/10.3176/earth.2014.07>
- Duarte, C. M., Delgado-Huertas, A., Anton, A., Carrillo-de-Albornoz, P., López-Sandoval, D. C., Agustí, S., Almahasheer, H., Marbá, N., Hendriks, I. E., Krause-Jensen, D., & Garcias-Bonet, N. (2018). Stable

- Isotope ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ,  $\delta^{18}\text{O}$ ,  $\delta\text{D}$ ) composition and nutrient concentration of Red Sea primary producers. *Frontiers in Marine Science*, 5, 1–12. <https://doi.org/10.3389/fmars.2018.00298>
- Duigan, C. A., & Birks, H. H. (2000). The late-glacial and early-Holocene palaeoecology of cladoceran microfossil assemblages at Krakenes, western Norway, with a quantitative reconstruction of temperature changes. *Journal of Paleolimnology*, 23(1), 67–76. <https://doi.org/10.1023/A:1008004513301>
- Ejankowski, W. (2008). Effect of waterlogging on regeneration in the dwarf birch (*Betula nana*). *Biologia*, 63(5), 670–676. <https://doi.org/10.2478/s11756-008-0126-8>
- Faegri, K. & Iversen, J. (1964). Textbook of Pollen Analysis. Hafner Publishing Company, New Yourk, 2 ed., 237 pp.
- Feurdean, A., Vanni re, B., Finsinger, W., Warren, D., Connor, S. C., Forrest, M., Liakka, J., Panait, A., Werner, C., Andri , M., Bobek, P., Carter, V. A., Davis, B., Diaconu, A., Dietze, E. et al. (2020). Fire hazard modulation by long-term dynamics in land cover and dominant forest type in eastern and central Europe. *Biogeosciences*, 17(5), 1213–1230. <https://doi.org/10.5194/bg-17-1213-2020>
- Fiamegos, Y., & de la Calle Gunti nas, M. B. (2018). Validation strategy for an ed-xrf method to determine trace elements in a wide range of organic and inorganic matrices based on fulfilment of performance criteria. *Spectrochimica Acta - Part B Atomic Spectroscopy*, 150, 59–66. <https://doi.org/10.1016/j.sab.2018.10.009>
- Ficken, K. J., Li, B., Swain, D. L., & Eglinton, G. (2000). An n-alkane proxy for the sedimentary input of submerged/floating freshwater aquatic macrophytes. *Organic Geochemistry*, 31(7–8), 745–749. [https://doi.org/10.1016/S0146-6380\(00\)00081-4](https://doi.org/10.1016/S0146-6380(00)00081-4)
- Firestone R. B., West, A., Kennett, J. P. , Becker, L., Bunch, T. E., Revay, Z. S., Schultz, P. H., Belgya, T., Kennett, D. J., Erlandson, J. M., Dickenson, O. J., Goodyear, A. C., Harris, R. S., Howard, G. A., Kloosterman, J. B., Lechler, P., Mayewski, P. A., Montgomery, J., Poreda, R., Darrah, T., Que Hee, S. S., Smith, A. R., Stich, A., Topping, W., Wittke, J. H. & Wolbach, W. S. (2007). Evidence for an extraterrestrial impact 12,900 years ago that contributed to the megafaunal extinctions and the Younger Dryas cooling. *Proceedings of the National Academy of Sciences*, 104(41), 16016-16021. <https://doi.org/10.1073/pnas.0706977104>
- Fott, J., Pra akov, M., Stuchlk, E., & Stuchlikov, Z. (1994). Acidification of lakes in  umava (Bohemia) and in the High Tatra Mountains (Slovakia). *Hydrobiologia*, 274(1–3), 37–47. <https://doi.org/10.1007/BF00014625>
- Frey, D. G. (1966). Contrasting Strategies of Gamogenesis in Northern and Southern Populations of Cladocera. *Ecology*, 63(1), 223–241. <https://doi.org/10.2307/1937046>
- Frolova, L., Nazarova, L., Pestryakova, L., & Herzsuh, U. (2014). Subfossil Cladocera from surface sediment in thermokarst lakes in northeastern Siberia, Russia, in relation to limnological and climatic variables. *Journal of Paleolimnology*, 52(1–2), 107–119. <https://doi.org/10.1007/s10933-014-9781-7>
- Gandouin, E., Rioual, P., Pailles, C., Brooks, S. J., Ponel, P., Guiter, F., Djamali, M., Andrieu-Ponel, V., Birks, H. J. B., Leydet, M., Belkacem, D., Haas, J. N., Van der Putten, N. & Beaulieu, J. L.(2016). Environmental and climate reconstruction of the late-glacial-Holocene transition from a lake sediment sequence in Aubrac, French Massif Central: Chironomid and diatom evidence. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 461, 292–309. <https://doi.org/10.1016/j.palaeo.2016.08.039>
- Gilbert, J. J. J. (1988). Suppression of rotifer population by *Daphnia* : A review of the evidence, the mechanisms, and the effects on zooplankton community structure. *Limnology and Oceanography*, 33(6), 1286–1303. <http://www.jstor.org/stable/10.2307/2837290>
- Goslar, T., Ralska-Jasiewiczowa, M., Van Geel, B., acka, B., Szeroczyńska, K., Chrst, L., & Walanus, A. (1999). Anthropogenic changes in the sediment composition of Lake Gosciaz (central Poland), during the last 330 yrs. *Journal of Paleolimnology*, 22(2), 171–185. <https://doi.org/10.1023/A:1008096032117>
- Goulden, C. E. (1971). Distribution Control of the of the Chydorid Abundance. *Limnology and Oceanography*, 16(2), 320–331.

- Guilizzoni, P., Lami, A., Marchetto, A., Jones, V., Manca, M., & Bettinetti, R. (2002). Palaeoproductivity and environmental changes during the Holocene in central Italy as recorded in two crater lakes (Albano and Nemi). *Quaternary International*, 88, 57–68. [https://doi.org/10.1016/S1040-6182\(01\)00073-8](https://doi.org/10.1016/S1040-6182(01)00073-8)
- Hann, B. J., & Zrum, L. (1997). Littoral microcrustaceans (Cladocera, Copepoda) in a prairie coastal wetland: Seasonal abundance and community structure. *Hydrobiologia*, 357, 37–52. <https://doi.org/10.1023/A:1003170416487>
- Hann, B. J., & Turner, M. A. (1999). Exploitation by microcrustacea of a new littoral habitat in an acidified lake. *Hydrobiologia*, 416, 65–75. <https://doi.org/10.1023/A:1003894511586>
- Hartz, I., & Milthers, V. (1901). Det senglacie ler i Allerød tegelværksgrav. *Meddelelser Danks Geologisk Foreningen*, 8, 31–60.
- Heiri, O., Brooks, S. J., Renssen, H., Bedford, A., Hazekamp, M., Ilyashuk, B., Jeffers, E. S., Lang, B., Kirilova, E., Kuiper, S., Millet, L., Samartin, S., Toth, M., Verbruggen, F., Watson, J. E., van Asch, N., Lammertsma, E., Amon, L., Birks, H. H., Birks, H. J. B. et al. (2014). Validation of climate model-inferred regional temperature change for late-glacial Europe. *Nature Communications*, 5, 1–7. <https://doi.org/10.1038/ncomms5914>
- Heiri, O., & Lotter, A. F. (2003). 9000 Years of chironomid assemblage dynamics in an Alpine lake: Long-term trends, sensitivity to disturbance, and resilience of the fauna. *Journal of Paleolimnology*, 30(3), 273–289. <https://doi.org/10.1023/A:1026036930059>
- Hepp, J., Wüthrich, L., Bromm, T., Bliedtner, M., Schäfer, I. K., Glaser, B., Rozanski, K., Sirocko, F., Zech, R., & Zech, M. (2019). How dry was the Younger Dryas? Evidence from a coupled  $\delta^2\text{H}$ - $\delta^{18}\text{O}$  biomarker paleohygrometer applied to the Gemündener Maar sediments, Western Eifel, Germany. *Climate of the Past*, 15(2), 713–733. <https://doi.org/10.5194/cp-15-713-2019>
- Hofmann, W. (1998). Cladocerans and chironomids as indicators of lake level changes in north temperate lakes. *Journal of Paleolimnology*, 19(1), 55–62. <https://doi.org/10.1023/A:1007999419765>
- Holand, S. M. (2008). DETRENDED CORRESPONDENCE ANALYSIS (DCA). *RStudio Tutorial*. Department of Geology, University of Georgia, Athens, 14 pp.
- Holliday, V., Surovell, T., & Johnson, E. (2016). A Blind Test of the Younger Dryas Impact Hypothesis. *PLoS ONE*, 11(7) 4–8. <https://doi.org/10.1371/journal.pone.0155470>
- Hošek, J., Pokorný, P., Kubov, V., Horá, I., Pavla, Ž., Kadlec, J., Rojik, F., Lisá, L., & Bučkuliaková, S. (2014). Late glacial climatic and environmental changes in eastern-central Europe : Correlation of multiple biotic and abiotic proxies from the Lake Švarcenberk, Czech Republic. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 396, 155–172 Contents 396, 155–172. <https://doi.org/10.1016/j.palaeo.2013.12.024>
- Hošek, J., Pokorný, P., Prach, J., Lisá, L., Grygar, T. M., Kněsl, I., & Trubač, J. (2017). Late Glacial erosion and pedogenesis dynamics: Evidence from high-resolution lacustrine archives and paleosols in south Bohemia (Czech Republic). *Catena*, 150, 261–278. <https://doi.org/10.1016/j.catena.2016.11.022>
- Hošek, J., Prach, J., Křížek, M., Šída, P., Moska, P., & Pokorný, P. (2019). Buried Late Weichselian thermokarst landscape discovered in the Czech Republic, central Europe. *Boreas*, 48(4), 988–1005. <https://doi.org/10.1111/bor.12404>
- Hošek, J., Prach, J., Šída, P., Houfková, P., Vondrák, D., Lisá, L., Pokorný, P., Chvojka, O., & Dohnal, J. (2013). Newly Discovered Late Glacial lakes in the Třeboň Region. *Geoscience Research Reports*. 46, 126-132. <https://doi.org/10.3140/zpravy.geol.2016.40>
- Hou, J., D'Andrea, W. J., & Huang, Y. (2008). Can sedimentary leaf waxes record D/H ratios of continental precipitation? Field, model, and experimental assessments. *Geochimica et Cosmochimica Acta*, 72(14), 3503–3517. <https://doi.org/10.1016/j.gca.2008.04.030>
- Houfková, P., Bešta, T., Bernardová, A., Vondrák, D., Pokorný, P., & Novák, J. (2006). Holocene climatic events linked to environmental changes at Lake Komořany Basin, Czech Republic. *Holocene*, 27(8), 1132–1145. <https://doi.org/10.1177/0959683616683250>

- Hrynowiecka, A., Żarski, M., Jakubowski, G., Nadachowski, A., Pawłowska, K., Pawłowski, D., Szymanek, M., & Nast, D. (2018). Eemian and Vistulian (Weichselian) paleoenvironmental changes: A multi-proxy study of sediments and mammal remains from the Ławy paleolake (Eastern Poland). *Quaternary International*, 467, 131–146. <https://doi.org/10.1016/j.quaint.2016.10.033>
- Huang, Y., Shuman, B., Wang, Y., & Webb, T. (2004). Hydrogen isotope ratios of individual lipids in lake sediments as novel tracers of climatic and environmental change: a surface sediment test. *Journal of Paleolimnology*, 31, 363–375. <https://doi.org/10.1023/B:JOPL.0000021855.80535.13>
- Ilyashuk, B., Gobet, E., Heiri, O., Lotter, A. F., van Leeuwen, J. F. N., van der Knaap, W. O., Ilyashuk, E., Oberli, F., & Ammann, B. (2009). Lateglacial environmental and climatic changes at the Maloja Pass, Central Swiss Alps, as recorded by chironomids and pollen. *Quaternary Science Reviews*, 28(13–14), 1340–1353. <https://doi.org/10.1016/j.quascirev.2009.01.007>
- Jankovská, V. (2006). Late Glacial and Holocene history of Plešné Lake and its surrounding landscape based on pollen and palaeoalgalogical analyses. *Biologia*, 61(20), 371–385. <https://doi.org/10.2478/s11756-007-0064-x>
- Johnsen, S., Clausen, H. B., Dansgaard, W., Gundestrup, N., Hansson, M., Jonsson, P., Steffensen, J. P. & Sveinbjørnsdóttir, A. E. (1992). A ‘deep’ ice core from East Greenland. *Meddelelser om Grønland, Geoscience*, 29, 22 pp.
- Jovanovska, E., Cvetkoska, A., Hauffe, T., Levkov, Z., Wagner, B., Sulpizio, R., Francke, A., Albrecht, Ch., & Wilke, T. (2016). Differential resilience of ancient sister lakes Ohrid and Prespa to environmental disturbances during the Late Pleistocene. *Biogeosciences*, 13(4), 1149–1161. <https://doi.org/10.5194/bg-13-1149-2016>
- Kletetschka, G., Vondrák, D., Hrubá, J., Procházka, V., Nabelek, L., Svitavská-Svobodová, H., Bobek, P., Hořícká, Z., Kadlec, J., Takač, M., & Stuchlík, E. (2018). Cosmic-Impact Event in Lake Sediments from Central Europe Postdates the Laacher See Eruption and Marks Onset of the Younger Dryas. *The Journal of Geology*, 126, 561–575. <https://doi.org/10.1086/699869>
- Korhola, A., Olander, H., & Blom, T. (2000). Cladoceran and chironomid assemblages as quantitative indicators of water depth in subarctic Fennoscandian lakes. *Journal of Paleolimnology*, 24(1), 43–54. <https://doi.org/10.1023/A:1008165732542>
- Korhola, A., Tikkanen, M., & Weckström, J. (2005). Quantification of Holocene lake-level changes in Finnish Lapland using a cladocera - Lake depth transfer model. *Journal of Paleolimnology*, 34(2), 175–190. <https://doi.org/10.1007/s10933-005-1839-0>
- Kulesza, P., Suchora, M., Pidek, I. A., & Alexandrowicz, W. P. (2011). Chronology and directions of Late Glacial paleoenvironmental changes: A multi-proxy study on sediments of Lake Słone (SE Poland). *Quaternary International*, 238(1–2), 89–106. <https://doi.org/10.1016/j.quaint.2010.12.024>
- Labaj, A. L., Korosi, J. B., Kurek, J., Jeziorski, A., Keller, W. B., & Smol, J. P. (2016). Response of *Bosmina* size structure to the acidification and recovery of lakes near Sudbury, Canada. *Journal of Limnology*, 75(2), 22–29. <https://doi.org/10.4081/jlimnol.2016.1183>
- Leng, M. J. (2010). A multiproxy palaeolimnological change, between c. 10 700 and 7200 years BP, at Holebudalen, southern Norway. *The Holocene*, 18(5), 805–817. <https://doi.org/10.1177/0959683608089217>
- Li, G., & Piper, D. J. W. (2015). The influence of meltwater on the Labrador Current in Heinrich event 1 and the Younger Dryas. *Quaternary Science Reviews*, 107, 129–137. <https://doi.org/10.1016/j.quascirev.2014.10.021>
- Locke, A., & Sprules, W. G. (2000). Effects of acidic pH and phytoplankton on survival and condition of *Bosmina longirostris* and *Daphnia pulex*. *Hydrobiologia*, 437, 187–196. <https://doi.org/10.1023/A:1026563109217>
- Lotter, A. F., Birks, H. J. B., Hofmann, W., & Marchetto, A. (1998). Modern diatom, cladocera, chironomid, and chrysophyte cyst assemblages as quantitative indicators for the reconstruction of past environmental

- conditions in the Alps. II. Nutrients. *Journal of Paleolimnology*, 19(4), 443–463.  
<https://doi.org/10.1023/A:1007994206432>
- Lowe, J. J., Rasmussen, S. O., Björck, S., Hoek, W. Z., Steffensen, J. P., Walker, M. J. C., & Yu, Z. C. (2008). Synchronisation of palaeoenvironmental events in the North Atlantic region during the Last Termination: a revised protocol recommended by the INTIMATE group. *Quaternary Science Reviews*, 27(1–2), 6–17.  
<https://doi.org/10.1016/j.quascirev.2007.09.016>
- Luoto, T. P., Nevalainen, L., Kultti, S., & Sarmaja-Korjonen, K. (2011). An evaluation of the influence of water depth and river inflow on quantitative Cladocera-based temperature and lake level inferences in a shallow boreal lake. *Hydrobiologia*, 676(1), 143–154. <https://doi.org/10.1007/s10750-011-0801-6>
- Magny, M., Guiot, J., & Schoellammer, P. (2001). Quantitative reconstruction of Younger Dryas to Mid-Holocene paleoclimates at Le Locle, Swiss Jura, using Pollen and lake-level data. *Quaternary Research*, 56(2), 170–180. <https://doi.org/10.1006/qres.2001.2257>
- Manca, M., Torretta, B., Comoli, P., Amsinck, S. L., & Jeppesen, E. (2007). Major changes in trophic dynamics in large, deep sub-alpine Lake Maggiore from 1940s to 2002: A high resolution comparative palaeoneolimnological study. *Freshwater Biology*, 52(11), 2256–2269. <https://doi.org/10.1111/j.1365-2427.2007.01827.x>
- Matějka, K. (2011). Rozbor průběhu počasí na Churáňově (Šumava) v období 1983-2010 a jeho možná interpretace z hlediska dynamiky ekosystémů. *www.infodatasys.cz*, 18 pp., <https://www.infodatasys.cz/climate/churanov1983-2010.pdf>
- Mentlík, P., Engel, Z., Braucher, R., Léanni, L., & Team, A. (2013). Chronology of the Late Weichselian glaciation in the Bohemian Forest in Central Europe. *Quaternary Science Reviews*, 65, 120–128. <https://doi.org/10.1016/j.quascirev.2013.01.020>
- Meyers, P. A. (1997). Organic geochemical proxies of paleoenvironmental, paleolimnologic and paleoclimatic processes. *Organic Geochemistry*, 27(5), 2013–2050.
- Meyers, P. A., & Lallier-vergès, E. (1999). Lacustrine sedimentary organic matter records of Late Quaternary paleoclimates. *Journal of Paleolimnology*, 21, 345–372.
- Middelburg, J. J., Weijden, C. H., & Woittiez, J. R. W. (1988). Chemical processes affecting the mobility of major, minor and trace elements during weathering of granitic rocks. *Chemical Geology*, 68(3–4), 253–273. [https://doi.org/10.1016/0009-2541\(88\)90025-3](https://doi.org/10.1016/0009-2541(88)90025-3)
- Millet, L., Rius, D., Galop, D., Heiri, O., & Brooks, S. J. (2012). Chironomid-based reconstruction of Lateglacial summer temperatures from the Ech palaeolake record (French western Pyrenees). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 315, 86–99. <https://doi.org/10.1016/j.palaeo.2011.11.014>
- Millet, L., Vanniere, B., Valérie, V., Magny, M., Disnar, J. R., Laggoun-Défarge, F., Walter-Simonnet, A. V., Bossuet, G., Ortu, E., & Beaulieu, J. (2007). Response of littoral chironomid communities and organic matter to late glacial lake — level, vegetation and climate changes at Lago dell'Accesa (Tuscany, Italy). *Journal of Paleolimnology*, 38, 525–539. <https://doi.org/10.1007/s10933-007-9088-z>
- Moreno, P. I., Kaplan, M. R., François, J. P., Moy, C. M., Stern, C. R., & Kubik, P. W. (2009). Renewed glacial activity during the Antarctic cold reversal and persistence of cold conditions until 11.5 ka in southwestern Patagonia. *Geology*, 37(4), 375–378. <https://doi.org/10.1130/g25399a.1>
- Müller, S., Tarasov, P. E., Andreev, A. A., & Diekmann, B. (2009). Late Glacial to Holocene environments in the present-day coldest region of the Northern Hemisphere inferred from a pollen record of Lake Billyakh, Verkhoyansk Mts, NE Siberia. *Climate of the Past*, 5(1), 73–84. <https://doi.org/10.5194/cp-5-73-2009>
- Nedbalová, L., Fott, J., Kohout, L., Vrba, J., Kopáček, J., Soldán, T., & Macek, M. (2006). Biological recovery of the Bohemian Forest lakes from acidification. *Biologia*, 61, 453–465. <https://doi.org/10.2478/s11756-007-0071-y>
- Neugebauer, I., Brauer, A., Dräger, N., Dulski, P., Wulf, S., Plessen, B., Mingram, J., Herzsuh, U., & Brande, A. (2012). A Younger Dryas varve chronology from the Rehwiese palaeolake record in NE-Germany.

- Nevalainen, L. (2008). Sexual reproduction in chydorid cladocerans (Anomopoda, Chydoridae) in southern Finland – implications for paleolimnology. Ph.D. thesis, 55 pp., Department of Geology, University of Helsinki
- Nevalainen, L., Helama, S., & Luoto, T. P. (2013). Hydroclimatic variations over the last millennium in eastern Finland disentangled by fossil Cladocera. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 378, 13–21. <https://doi.org/10.1016/j.palaeo.2013.03.016>
- Nevalainen, L., Luoto, T. P., Kultti, S., & Sarmaja-Korjonen, K. (2012). Do subfossil Cladocera and chydorid ephippia disentangle Holocene climate trends? *Holocene*, 22(3), 291–299. <https://doi.org/10.1177/0959683611423691>
- Nevalainen, L., Rantala, M. V., & Luoto, T. P. (2015). Sedimentary cladoceran assemblages and their functional attributes record late Holocene climate variability in southern Finland. *Journal of Paleolimnology*, 54(2–3), 239–252. <https://doi.org/10.1007/s10933-015-9849-z>
- Niska, M., & Mirosław-Grabowska, J. (2014). Eemian environmental changes recorded in lake deposits from Rzecino (NW Poland): Cladocera, isotopic and selected geochemical data. *Journal of Paleolimnology*, 53(1), 89–105. <https://doi.org/10.1007/s10933-014-9810-6>
- Nyman, M., Weckström, J., & Korhola, A. (2008). Chironomid response to environmental drivers during the Holocene in a shallow treeline lake in northwestern Fennoscandia. *Holocene*, 18(2), 215–227. <https://doi.org/10.1177/0959683607086760>
- Olesen, J. (1998). A phylogenetic analysis of the Conchostraca and Cladocera (Crustacea, Branchiopoda, Diplostraca). *Zoological Journal of the Linnean Society*, 122, 491–536.
- Pawłowski, D. (2017). The usefulness of subfossil Cladocera remains in Younger Dryas climatic reconstructions in central Poland. *Acta Geologica Polonica*, 67(4), 567–584. <https://doi.org/10.1515/aggp-2017-0026>
- Pawłowski, D., Kloss, M., Obremska, M., Szymanowski, M., & Zurek, S. (2012). Evolution of small valley mire in central Poland as a result of hydroclimatic oscillations. *Geochronometria*, 39(2), 133–148. <https://doi.org/10.2478/s13386-012-0004-6>
- Pedziszewska, A., Tylmann, W., Witak, M., Piotrowska, N., Maciejewska, E., & Latałowa, M. (2015). Holocene environmental changes reflected by pollen, diatoms, and geochemistry of annually laminated sediments of Lake Suminko in the Kashubian Lake District (N Poland). *Review of Palaeobotany and Palynology*, 216, 55–75. <https://doi.org/10.1016/j.revpalbo.2015.01.008>
- Petera-Zganiacz, J., Borówka, R. K., Forsytek, J., Miotk-Szpiganowicz, G., & Pawłowski, D. (2013). Palaeogeographical significance of the Eemian biogenic sediments at the bór site (Warta river valley, central Poland). *Geological Quarterly*, 57(4), 687–700. <https://doi.org/10.7306/gq.1122>
- Petera-Zganiacz, J., Dzieduszyńska, D. A., Twardy, J., Pawłowski, D., Płóciennik, M., Lutyńska, M., & Kittel, P. (2015). Younger Dryas flood events: A case study from the middle Warta River valley (Central Poland). *Quaternary International*, 386, 55–69. <https://doi.org/10.1016/j.quaint.2014.09.074>
- Peyron, O., & Be, C. (2005). Late-Glacial climatic changes in Eastern France (Lake Lautrey) from pollen, lake-levels, and chironomids, 64, 197–211. <https://doi.org/10.1016/j.yqres.2005.01.006>
- Pigati, J. S., Latorre, C., Rech, J. A., Betancourt, J. L., Martínez, K. E., & Budahn, J. R. (2012). Accumulation of impact markers in desert wetlands and implications for the Younger Dryas impact hypothesis. *Proceedings of the National Academy of Sciences*, 109(19), 7208–7212. <https://doi.org/10.1073/pnas.1200296109>
- Piscia, R., Guilizzoni, P., Fontaneto, D., Vignati, D. A. L., Appleby, P. G., & Manca, M. (2012). Dynamics of rotifer and cladoceran resting stages during copper pollution and recovery in a subalpine lake. *Annales de Limnologie - International Journal of Limnology*, 48(2), 151–160. <https://doi.org/10.1051/limn/2012006>
- Płóciennik, M., Self, A., Birks, H. J. B., & Brooks, S. J. (2011). Chironomidae (Insecta: Diptera) succession in Zabieniec bog and its palaeo-lake (central Poland) through the Late Weichselian and Holocene.

*Palaeogeography, Palaeoclimatology, Palaeoecology*, 307(1–4), 150–167.  
<https://doi.org/10.1016/j.palaeo.2011.05.010>

- Pražáková, M., Veselý, J., Fott, J., Majer, V., & Kopáček, J. (2006). The long-term succession of cladoceran fauna and palaeoclimate forcing : A 14, 600 – year record from Plešné Lake, the Bohemian Forest. *Biologia*, 61(20), 387-399. <https://doi.org/10.2478/s11756-007-0072-x>
- Rautio, M. (2007). PALEOLIMNOLOGY/Cladocera. In: Elias, S.A. (ed) *Encyclopedia of Quaternary Science*, pp. 2029–2038. Elsevier Science, ISBN 978-0-444-52747-9.  
<https://doi.org/10.1016/B0-44-452747-8/00246-5>
- Renssen, H., van Geel, B., van der Plicht, J. & Magny, M. (2000). Reduced solar activity as a trigger for the start of the Younger Dryas ? *Quaternary International*, 68, 373-383.  
[https://doi.org/10.1016/S1040-6182\(00\)00060-4](https://doi.org/10.1016/S1040-6182(00)00060-4)
- Renssen, H., Mairesse, A., Goosse, H., Mathiot, P., Heiri, O., Roche, D. M., Nisancioglu, K. H., & Valdes, P. J., (2015). Multiple causes of the Younger Dryas cold period. *Nature Geoscience*, 8, 946–949.  
<https://doi.org/10.1038/NGEO2557>
- Renssen, H., Isarin, R. F. B., Vandenberghe, J., & Lautenschlager, M. (2000). Permafrost as a critical factor in paleoclimate modelling : the Younger Dryas case in Europe. *Earth and Planetary Science Letters*, 176, 1–5. [https://doi.org/10.1016/S0012-821X\(99\)00322-2](https://doi.org/10.1016/S0012-821X(99)00322-2)
- Sacherová, V., Kršková, R., Stuchlík, E., Hořická, Z., Hudec, I., & Fott, J. (2006). Long-term change of the littoral Cladocera in the Tatra Mountain lakes through a major acidification event. *Biologia*, 61, 109–119.  
<https://doi.org/10.2478/s11756-006-0124-7>
- Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R., McInerney, F. A., van der Meer, M. T. J., Polissar, P., Robins, R. J., Sachs, J. P., Schmidt, H., Sessions, A. L., White, J. W. C., West, J. B., & Kahmen, A. (2012). Molecular paleohydrology: Interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annual Review of Earth and Planetary Sciences*, 40, 221–249. <https://doi.org/10.1146/annurev-earth-042711-105535>
- Samartin, S., Heiri, O., Vescovi, E., Brooks, S. J., & Tinner, W. (2012). Lateglacial and early Holocene summer temperatures in the southern Swiss Alps reconstructed using fossil chironomids. *Journal of Quaternary Science*, 27(3), 279–289. <https://doi.org/10.1002/jqs.1542>
- Sarmaja-Korjonen, K. (2002). Multi-proxy data from Kaksoislammi Lake in Finland: Dramatic changes in the late Holocene cladoceran assemblages. *Journal of Paleolimnology*, 28(3), 287–296.  
<https://doi.org/10.1023/A:1021611225821>
- Sarmaja-Korjonen, K. (2004). Chydorid ephippia as indicators of past environmental changes - A new method. *Hydrobiologia*, 526, 129–136. <https://doi.org/10.1023/B:HYDR.0000041595.08121.ab>
- Sarmaja-Korjonen, K., & Alhonen, P. (1999). Cladoceran and diatom evidence of lake-level fluctuations from a Finnish lake and the effect of aquatic-moss layers on microfossil assemblages. *Journal of Paleolimnology*, 22(3), 277–290. <https://doi.org/10.1023/A:1008061702639>
- Sarmaja-Korjonen, K., Nyman, M., Kultti, S., & Väiliranta, M. (2006). Palaeolimnological development of Lake Njargajavri, northern Finnish Lapland, in a changing Holocene climate and environment. *Journal of Paleolimnology*, 35(1), 65–81. <https://doi.org/10.1007/s10933-005-7337-6>
- Sarmaja-Korjonen, K., & Seppä, H. (2007). Abrupt and consistent responses of aquatic and terrestrial ecosystems to the 8200 cal. yr cold event: A lacustrine record from Lake Arapisto, Finland. *Holocene*, 17(4), 457–467. <https://doi.org/10.1177/0959683607077020>
- Sernander, R. (1908). On the evidence of post-glacial changes of climate furnished by the peat mosses of northern Europe. *Geologiska Föreningens i Stockholm Förhållinger*, 30, 365-478.
- Shuman, B., Bartlein, P. J., & Webb, T. (2005). The magnitudes of millennial- and orbital-scale climatic change in eastern North America during the Late Quaternary. *Quaternary Science Reviews*, 24, 2194–2206.  
<https://doi.org/10.1016/j.quascirev.2005.03.018>

- Siebel, W., Shang, C. K., Reitter, E., Rohrmüller, J., & Breiter, K. (2008). Two distinctive granite suites in the SW Bohemian Massif and their record of emplacement: constraints from geochemistry and zircon  $^{207}\text{Pb}/^{206}\text{Pb}$  chronology. *Journal of Petrology*, 49(10), 1853–1872. <https://doi.org/10.1093/petrology/egn049>
- Siitonen, S., Väiliranta, M., Weckström, J., Juutinen, S., & Korhola, A. (2011). Comparison of Cladocera-based water-depth reconstruction against other types of proxy data in Finnish Lapland. *Hydrobiologia*, 676(1), 155–172. <https://doi.org/10.1007/s10750-011-0885-z>
- Skurčáková, A. (2018). Paleoenvironmentální rekonstrukce mladšího dryasu na základě fosilních pakomárů, Diplomová práce, 84 str., Přírodovědecká fakulta, Univerzita Karlova, Praha
- Słowiński, M., Marcisz, K., Plóciennik, M., Obremska, M., Pawłowski, D., Okupny, D., Okupny, D., Słowińska, S., Borówka, R., Kittel, P., Forysiak, J., Michczyńska, D. J., & Lamentowicz, M. (2016). Drought as a stress driver of ecological changes in peatland - A palaeoecological study of peatland development between 3500 BCE and 200 BCE in central Poland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 461, 272–291. <https://doi.org/10.1016/j.palaeo.2016.08.038>
- Smol, J. P., Birks, H. J. B., & Last, W. M. (eds.). (2001). Tracking environmental change using lake sediments. Volume 4: Zoological indicators. Springer, Dordrecht, 218 pp. <https://doi.org/10.1007/0-306-47671-1>
- Stankevica, K., Kalnina, L., Klavins, M., Cerina, A., Ustupe, L., & Kaup, E. (2015). Reconstruction of the Holocene palaeoenvironmental conditions accordingly to the multiproxy sedimentary records from Lake Pilvelis, Latvia. *Quaternary International*, 386, 102–115. <https://doi.org/10.1016/j.quaint.2015.02.031>
- Stefanova, I., Ognjanova-Rumenova, N., Hofmann, W., & Ammann, B. (2003). Late Glacial and Holocene environmental history of the Pirin Mountains (SW Bulgaria): A paleolimnological study of Lake Dalgoto (2310 m). *Journal of Paleolimnology*, 30(1), 95–111. <https://doi.org/10.1023/A:1024746818627>
- Stocker, T. F., & Johnsen, S. J. (2003). A minimum thermodynamic model for the bipolar seesaw. *Paleoceanography*, 18(4), 1–9. <https://doi.org/10.1029/2003PA000920>
- Sweetman, J. N., & Smol, J. P. (2006). Patterns in the distribution of cladocerans (Crustacea: Branchiopoda) in lakes across a north-south transect in Alaska, USA. *Hydrobiologia*, 553(1), 277–291. <https://doi.org/10.1007/s10750-005-1333-8>
- Szeroczyńska, K., & Sarmaja-Korjonen, K. (2007). Atlas of subfossil cladocera from central and northern Europe. Friends of Lower Vistula Society, Swiecie, 84 pp. ISBN 978-83-924919-6-5.
- Szeroczyńska, K. (1986). Cladocera jako wskaźnik ekologiczny w Późnoczwartorzędowych osadach jeziornych Polski Północnej. *Paleolimnologia*, 30, 3-69.
- Szeroczyńska, K. (1998). Palaeolimnological investigations in Poland based on Cladocera (Crustacea). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 140(1–4), 335–345. [https://doi.org/10.1016/S0031-0182\(98\)00028-5](https://doi.org/10.1016/S0031-0182(98)00028-5)
- Tanoi, T., Kawachi, M., & Watanabe, M. M. (2011). Effects of carbon source on growth and morphology of *Botryococcus braunii*. *Journal of Applied Phycology*, 23(1), 25–33. <https://doi.org/10.1007/s10811-010-9528-4>
- Taylor, D. J., Crease, T. J., & Brown, W. M. (1999). Phylogenetic evidence for a single long-lived clade of crustacean cyclic parthenogens and its implications for the evolution of sex. *Proceedings of the Royal Society B: Biological Sciences*, 266(1421), 791–797. <https://doi.org/10.1098/rspb.1999.0707>
- Teranes, J. L., & Bernasconi, S. M. (2005). Factors controlling  $\delta^{13}\text{C}$  values of sedimentary carbon in hypertrophic Baldeggersee, Switzerland, and implications for interpreting isotope excursions in lake sedimentary records. *Limnology and Oceanography*, 50(3), 914–922. <https://doi.org/10.4319/lo.2005.50.3.0914>
- Texlová, E. (2010). Vodní plošnice ( Heteroptera : Nepomorpha, Gerromorpha ) šumavských jezer, Diplomová práce, 93 str., Pedagogická fakulta, Jihočeská univerzita v Českých Budějovicích

- Thevenon, F., Adatte, T., Spangenberg, J. E., & Anselmetti, F. S. (2012). Elemental (C/N ratios) and isotopic ( $\delta^{15}\text{N}_{\text{org}}$ ,  $\delta^{13}\text{C}_{\text{org}}$ ) compositions of sedimentary organic matter from a high-altitude mountain lake (Meidsee, 2661 m a.s.l., Switzerland): Implications for Lateglacial and Holocene Alpine landscape evolution. *The Holocene*, 22(10), 1135–1142. <https://doi.org/10.1177/0959683612441841>
- Tinner, W., & Kaltenrieder, P. (2005). Rapid responses of high-mountain vegetation to early Holocene environmental changes in the Swiss Alps. *Journal of Ecology*, 93(5), 936–947. <https://doi.org/10.1111/j.1365-2745.2005.01023.x>
- Torres, I. C., Inglett, P. W., Brenner, M., & Kenney, W. F. (2012). Stable isotope ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) values of sediment organic matter in subtropical lakes of different trophic status. *Journal of Paleolimnology*, 47, 693–706. <https://doi.org/10.1007/s10933-012-9593-6>
- Tóth, A., Horváth, Z., Vad, C. F., Zsuga, K., Nagy, S. A., & Boros, E. (2014). Zooplankton of the European soda pans: Fauna and conservation of a unique habitat type. *International Review of Hydrobiology*, 99(3), 255–276. <https://doi.org/10.1002/iroh.201301646>
- Turek, J., Fluksová, H., Hejzlar, J., Kopáček, J., & Porcal, P. (2014). Modelling air temperature in catchments of Čertovo and Plešné lakes in the Bohemian Forest back to 1781. *Silva Gabreta*, 20(1), 1–24.
- Vočadlova, K., Petr, L., Pavla, Z., & Ara, K. L. (2015). The Lateglacial and Holocene in Central Europe: a multi-proxy environmental record from the Bohemian Forest, Czech Republic. *Boreas*, 44(4), 769–784. <https://doi.org/10.1111/bor.12126>
- von Grafenstein, U., Eicher, U., Erlenkeuser, H., Ruch, P., Schwander, J., & Ammann, B. (2000). Isotope signature of the Younger Dryas and two minor oscillations at Gerzensee (Switzerland): palaeoclimatic and palaeolimnologic interpretation based on bulk and biogenic carbonates. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 159, 215–229.
- Vrba, J., Kopáček, J., Fott, J., Kohout, L., Nedbalová, L., Pražáková, M., Soldán, T., & Schaumburg, J. (2003). Long-term studies (1871–2000) on acidification and recovery of lakes in the Bohemian Forest (central Europe). *Science of the Total Environment*, 310(1–3), 73–85. [https://doi.org/10.1016/S0048-9697\(02\)00624-1](https://doi.org/10.1016/S0048-9697(02)00624-1)
- Wachecka-Kotkowska, L., Krzyszkowski, D., Malkiewicz, M., Mirosław-Grabowska, J., Niska, M., Krzysińska, J., Myśkow, E., Raczyk, J., Wieczorek, D., Stoiński, A., & Rządziejewicz, M. (2018). An attempt to reconstruct the late Saalian to Plenivistulian (MIS6–MIS3) natural lake environment from the “Parchliny 2014” section, central Poland. *Quaternary International*, 467, 5–25. <https://doi.org/10.1016/j.quaint.2016.06.013>
- Walker, M. J. C. (1995). Climatic Changes During the Last Glacial/Interglacial Transition. *Quaternary International*, 28, 63–76. [https://doi.org/10.1016/1040-6182\(95\)00030-M](https://doi.org/10.1016/1040-6182(95)00030-M)
- Wunsch, C. (2005). Abrupt Climate Change. An Alternative View. *Quaternary Research*, 65, 191–203. <https://doi.org/10.1016/j.yqres.2005.10.006>
- Yiou, P., & Jouzel, J. (1995). Rapid oscillations in Vostok and GRIP Ice Cores. *Geophysical Research Letters*, 22(16), 2179–2182. <https://doi.org/10.1029/95GL02014>
- Yu, N., Li, E., Feng, D., Xiao, B., Wei, C., Zhang, M., & Chen, L. (2014). Correlations between zooplankton assemblages and environmental factors in the downtown rivers of Shanghai, China. *Chinese Journal of Oceanology and Limnology*, 32(6), 1352–1363. <https://doi.org/10.1007/s00343-014-3325-x>
- Zachowicz, J. (1982). The late-Glacial and Holocene vegetational history of the Zulawy Region, N. Poland. A biostratigraphic study of Lake Druzno sediments (Vistula Lagoon). *Acta Palaeobotanica*, 22(1), 141–161.
- Zawiska, I., Słowiński, M., Correa-Metrio, A., Obremaska, M., Luoto, T., Nevalainen, L., Woszczyk, M., & Milecka, K. (2014). The response of a shallow lake and its catchment to Late Glacial climate changes - A case study from eastern Poland. *Catena*, 126, 1–10. <https://doi.org/10.1016/j.catena.2014.10.007>
- Zawisza, E., & Szeroczyńska, K. (2007). The development history of Wigry lake as shown by subfossil Cladocera. *Geochronometria*, 27(1), 67–74. <https://doi.org/10.2478/v10003-007-0021-2>

**Internetové stránky:**

Český hydrometeorologický ústav: <http://portal.chmi.cz>

Plán péče NPR Černé a Čertovo jezero: <https://drusop.nature.cz>

**Programy:**

Bennet, K. D. (1995–2007). Psimpoll 4.27

Grimm, E. C. (1991–2015). Tilia 2.6.1

RStudio 1.3.959 (2009-2020). RStudio, Inc.