

Hydrogen-Bonded Complexes of Phenylacetylene with Water, Methanol, Ammonia and Methylamine. The Origin of Methyl Group Induced Hydrogen Bond Switching.

Robert Sedlak,[†] Pavel Hobza,^{†,‡,*} and G. Naresh Patwari^{¶,*}

[†] Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic and Center for Biomolecules and Complex Molecular Systems, Flemingovo nám. 2, 166 10 Prague 6, Czech Republic

[‡] Department of Physical Chemistry, Palacký University, 771 46 Olomouc, Czech Republic

[¶] Department of Chemistry, Indian Institute of Technology Bombay, Powai, Mumbai 400076 India.

* Authors to whom correspondence should be addressed. E-mail: pavel.hobza@uochb.cas.cz (P.H.); naresh@chem.iitb.ac.in (G.N.P.)

Abstract

The infrared spectra in the acetylenic C–H stretching region for the complexes of phenylacetylene with water, methanol, and ammonia, methylamine are indicative of change in the intermolecular structure upon substitution with a methyl group. High-level ab initio calculations at CCSD(T)/aug-cc-pVDZ level indicate that the observed complexes of water and ammonia are energetically the most favored structures and electrostatics play dominant role in stabilizing these structures. The ability of the π electron density of the benzene ring to offer larger cross-section for the interaction and the increased polarizability of the O–H and N–H groups in methanol and methylamine favor the formation of π hydrogen-bonded complexes, in which dispersion is the dominate force. Further, the observed phenylacetylene-methylamine complex can be tentatively assigned to a kinetically trapped higher energy structure. The observed methyl group induced hydrogen bond switching in the phenylacetylene complexes can be attributed to the switching of the dominant interaction from electrostatic to dispersion.

Introduction

Phenylacetylene is perhaps the simplest multifunctional molecule to investigate competitive hydrogen bonding. It has three hydrogen bonding sites in the form of benzene ring and the acetylenic $\text{C}\equiv\text{C}$ bond, which can act as π acceptors, and an activated acetylenic C–H group, which can act as a σ donor. Further, with the absence of any strongly acidic/basic functional groups, the hierarchy of hydrogen sites cannot be

determined on the basis of Etter and Legon-Millen rules.^{1,2} One of the major challenges that needs to be addressed in hydrogen bonding is to know, a priori, how the individual functional groups in multifunctional molecules will behave when they are made to interact with suitable hydrogen bonding partners. In multifunctional molecules the exact hydrogen bonding pattern will be a result of competition between various possibilities. Towards the goal of comprehending the hydrogen bonding behavior of multifunctional molecules, Patwari and coworkers investigated hydrogen-bonded complexes of phenylacetylene with various solvent molecules such as water, methanol, ammonia, methylamine and other alcohols and amines.³⁻⁵ The hydrogen-bonded complexes of phenylacetylene form a wide variety of intermolecular structures, which stem out of subtle balance of intermolecular forces in various possible intermolecular structures.³⁻⁵ For instance, phenylacetylene forms a quasi-planar cyclic complex with water incorporating O–H $\cdots\pi$ and C–H \cdots O hydrogen bonds.^{3,4} In this case one of the O–H groups of water molecule interacts with the π electron density of the C \equiv C bond, while the C–H group of the benzene ring in the ortho position is hydrogen-bonded to the oxygen atom of the water molecule. Structure of phenylacetylene-water complex thus is different from both the benzene-water and acetylene-water complexes,^{6,7} even though phenylacetylene incorporates the features of both benzene and acetylene. On the other hand, the phenylacetylene-methanol complex is characterized by the presence of single O–H $\cdots\pi$ hydrogen bond, wherein the O–H group of methanol interacts with the π electron density of the benzene ring, similar to benzene-methanol complex.⁴ Phenylacetylene forms a linear C–H \cdots N “ σ ” hydrogen-bonded complex with ammonia,⁵ which is similar to acetylene-ammonia complex,⁸ while the phenylacetylene-methylamine

complex is characterized by the presence of N–H $\cdots\pi$ hydrogen bond. In this case the N–H group of methylamine interacts with the π electron density of the benzene ring.⁵ Such differences in the intermolecular structures of hydrogen-bonded complexes with water, methanol and ammonia, methylamine involving benzene derivatives have not been reported in the literature, prior to the complexes of phenylacetylene. These results illustrate that, in the case of interaction with multifunctional molecules even minimal changes in the interacting partner, such as substitution by a ubiquitous methyl group, can result in dramatic change in the intermolecular structure. The change in the intermolecular structure with the substitution of methyl group can be perceived as methyl group induced hydrogen bond switching. In this article we address the underlying factors that influence the hydrogen bond switching observed in the complexes of phenylacetylene.

Methodology

The geometry optimizations were carried out at MP2(FC)/aug-cc-pVDZ level of theory and in each case frequency calculations followed to ascertain the nature of the minima obtained. Single point calculations at CCSD(T)/aug-cc-pVDZ level were carried out on the MP2 level optimized structures. The stabilization energies were corrected for zero point energy (ZPE) and basis set superposition error (BSSE). Thermochemical analysis, based on rigid rotor – harmonic oscillator – ideal gas approximations was also carried out using the vibrational frequency data obtained at MP2(FC)/aug-cc-pVDZ level of theory. Further, DFT-SAPT (Symmetry Adapted Perturbation Theory) calculations were performed with the aug-cc-pVDZ basis set.⁹ This method allows for the separation

of interaction energies into physically well defined components, such as those arising from electrostatic, induction, dispersion and exchange. The DFT-SAPT interaction energy (E^{int}) is given as

$$E^{\text{int}} = E_1^{\text{Pol}} + E_1^{\text{Ex}} + E_2^{\text{Ind}} + E_2^{\text{Ex-Ind}} + E_2^{\text{D}} + E_2^{\text{Ex-D}} + \delta\text{HF} \quad (1)$$

Equation 1 describes the electrostatic, exchange-repulsion, induction, exchange-induction, dispersion and exchange-dispersion terms, while the last term is a Hartree-Fock correction for higher-order contributions to the interaction energy. In the present analysis the exchange-induction and exchange-dispersion terms will be included to the parent induction and dispersion terms. All calculations mentioned above were performed using Gaussian-03 and Molpro suit of programs.^{10,11}

Results and Discussion

To begin with, the IR spectra of phenylacetylene and its complexes in the acetylenic C–H stretching region are reviewed to understand the structural assignment. The infrared spectra in the acetylenic C–H stretching region for phenylacetylene (PHA) along with its five complexes with argon (Ar), water (H₂O), methanol (MeOH), ammonia (NH₃), and methylamine (MeNH₂) are depicted in Figure 1. These spectra were recorded using IR-UV double resonance spectroscopic method using either fluorescence or ion detection techniques.³⁻⁵ The IR spectrum of PHA (Trace A) shows the presence of two intense transitions at 3325 and 3343 cm⁻¹, which were assigned to be originating due to Fermi resonance coupling between the acetylenic C–H stretching vibration and a

combination band of comprising of one quantum of $\text{C}\equiv\text{C}$ stretching and two quanta of $\text{C}\equiv\text{C}-\text{H}$ out-of-plane bend.¹² A two state deperturbation analysis places the unperturbed acetylenic $\text{C}-\text{H}$ oscillator at 3334 cm^{-1} with the magnitude of coupling constant to be 9 cm^{-1} .¹³ In the case of PHA complexes any interaction that will perturb either the acetylenic $\text{C}-\text{H}$ oscillator or the $\text{C}\equiv\text{C}$ oscillator or both will lead to disappearance of Fermi resonance transitions. However, the perturbation should be about the order of the coupling constant of 9 cm^{-1} or more, in order completely remove the Fermi mixing. The Fermi resonance transitions of PHA moiety, therefore, can be used a spectroscopic tool to probe the interactions present in various PHA complexes. Figure 1B depicts the IR spectrum of PHA-argon complex, which is almost identical to that of bare PHA, within the experimental uncertainty of $\pm 1\text{ cm}^{-1}$.¹³ This implies that the binding of Ar to PHA does not perturb either the $\text{C}-\text{H}$ or the $\text{C}\equiv\text{C}$ oscillators. It can therefore be inferred that Ar atom is bound to the π electron density of the benzene ring in PHA.¹³ This inference is substantiated by the structure of PHA-Ar complex determined using high-resolution REMPI spectrum for the $S_1 \leftarrow S_0$ electronic transition and microwave spectroscopy.¹⁴

The IR spectrum of PHA- H_2O complex, depicted in Figure 1C, consists of a single transition at 3331 cm^{-1} , which has been assigned to the acetylenic $\text{C}-\text{H}$ stretching vibration. The acetylenic $\text{C}-\text{H}$ stretching frequency shifts marginally (about 3 cm^{-1}) upon interaction with water, coupled with the loss of Fermi resonance coupling. This implies that the H_2O molecule interacts with the π electron density of the acetylenic $\text{C}\equiv\text{C}$ bond.³ Surprisingly, in the case of PHA-MeOH (Figure 1D) complex two transitions appear at 3323 and 3334 cm^{-1} , albeit the differences in the band positions and their intensities,

relative to bare PHA, these transitions have been assigned to the Fermi resonance bands.⁴ In combination with the shift in the electronic transition for the $S_1 \leftarrow S_0$ relative to bare PHA, shift in the O–H stretching frequency of the MeOH moiety and the appearance of Fermi resonance transitions in the IR spectrum, it was assigned that MeOH interacts with the π electron density of the benzene ring in PHA resulting in formation of a O–H $\cdots\pi$ hydrogen-bonded complex.⁴ The positions and the intensities of the Fermi resonance bands depend on the positions of the zero-order (unperturbed) oscillators and the coupling strength. The interaction of methanolic O–H group with the π electron density of the benzene ring in PHA is expected to affect, marginally, both the zero-order positions and the coupling strength.

The assignment of the PHA-NH₃ complex to a C–H \cdots N hydrogen-bonded complex is rather straightforward due to the fact that the acetylenic C–H stretching vibration shifts by 103 cm⁻¹ to a lower frequency (see Figure 1E). Surprisingly, the IR spectrum of PHA-MeNH₂, depicted in Figure 1F, shows a single transition at 3333 cm⁻¹. The acetylenic C–H stretching frequency shifts by -1 cm⁻¹ upon interaction with MeNH₂, therefore completely rules out the possibility of formation of a C–H \cdots N hydrogen-bonded complex, similar to PHA-NH₃ complex. It is well known that the substitution of alkyl groups on NH₃ increases the basicity, which in-turn is expected to enhance the C–H \cdots N hydrogen-bonded interaction. Such a consideration would indicate that the acetylenic C–H stretching frequency of PHA should be further lowered upon interaction with MeNH₂, relative to the NH₃ complex. The analysis of the IR spectrum along with the electronic spectrum leads to the assignment of a structure which is characterized by the formation of N–H $\cdots\pi$ (benzene π) hydrogen bonding along with a with a peripheral

interaction between the methyl C–H group and the acetylenic C≡C bond.⁵ One of the interesting observations about this complex is that the MeNH₂ behaves as hydrogen bond donor, which is rather surprising considering the fact that alkyl amines are known to be excellent hydrogen bond acceptors and poor hydrogen bond donors. Additionally, examples in which the N–H group of aliphatic amines acts as a hydrogen bond donor are extremely sparse, to the extent of being non-existent in the gas-phase complexes.¹⁵

The most interesting aspect, which has been observed for the PHA complexes, is the difference in the intermolecular structure between H₂O and MeOH complexes, and similarly between NH₃ and MeNH₂ complexes. These observations can be summed up as methyl group induced hydrogen bond switching. To understand the origin of the observed hydrogen bond switching upon substitution with the methyl group, high-level ab initio calculations were carried out. To begin with geometries of the monomers and the complexes were optimized using MP2/aug-cc-pVDZ level of theory. Coincidentally, three minima were found for each set of complexes, structures of which are depicted in Figure 2. The first minimum in each case (**PWS1**, **PMS1**, **PAS1**, and **PMAS1**) is a linear C–H···O/N hydrogen-bonded complex, wherein the acetylenic C–H group acts as hydrogen bond donor to the O/N atom of the acceptor, structures similar to acetylene-H₂O and acetylene-NH₃ complexes.^{7,8} The second set of complexes are the π hydrogen-bonded complexes (**PWS2**, **PMS2**, **PAS2**, and **PMAS2**). In these cases the π electron density on the benzene ring of phenylacetylene is the hydrogen bond acceptor for the O–H/N–H groups of the interacting partner, similar to H₂O and NH₃ complexes of benzene.^{6,16} In the third set PHA forms a cyclic complexes with H₂O, NH₃ and MeNH₂, (**PWS3**, **PAS3**, and **PMAS3**). In these complexes both PHA and the interacting molecule act as hydrogen

bond donor and acceptor simultaneously, with the O–H/N–H group of the interacting molecule hydrogen-bonded to the π electron density of the C \equiv C of the acetylenic moiety in PHA. Further, the C–H group of the benzene ring in the ortho position is hydrogen-bonded to O/N of the accepting base. However, the interaction of MeOH with the π electron density of the C \equiv C is distinctly different from the other three complexes. The MeOH complex, **PMS3**, can be characterized by the presence of O–H $\cdots\pi$ (acetylene π) and of C–H $\cdots\pi$ (benzene π) hydrogen bonds. Several attempts were made to obtain the cyclic structure starting from modified initial geometries; however, in all the cases, the calculations converged on to the **PMS3** structure.

Based on the IR spectra in the acetylenic C–H stretching region (Figure 1) and other spectroscopic inputs,¹⁷ the structures of H₂O, MeOH, NH₃ and MeNH₂ complexes were assigned to **PWS3**, **PMS2**, **PAS1** and **PMAS2**, respectively.³⁻⁵ These structural assignments were supplemented by stabilization energies calculated at MP2/aug-cc-pVDZ with ZPE and 50% BSSE correction.³⁻⁵ The stabilization energies corrected for ZPE and 100% BSSE for all the complexes are listed in Table 1. However, with 100% BSSE correction only the observed structures of PHA-MeOH (**PMS2**) and PHA-NH₃ (**PAS1**) were found to be global minima for the respective complexes, while the observed structures of PHA-H₂O (**PWS3**) and PHA-MeNH₂ (**PMAS2**) correspond to higher energy local minima (see Table 1). The MP2/aug-cc-pVDZ level is known to provide accurate geometries of isolated systems as well as molecular complexes.¹⁸ On the other hand, it is also well known that MP2 level calculations overestimate the dispersion energies and single point calculations at CCSD(T) level would provide a more accurate description.¹⁹ Table 1 also lists ZPE and 100% BSSE corrected stabilization energies calculated at

CCSD(T) level. The difference in the stabilization energies calculated at the MP2 and the CCSD(T) levels are only marginal for the C–H \cdots O/N hydrogen-bonded complexes. However, the differences are substantial for rest of the complexes, which are characterized by the presence of π hydrogen bonding. This substantiates our earlier statement that MP2 level calculations overestimate the dispersion energies. At the CCSD(T) level the experimentally observed structure is the global minimum for the H₂O (**PWS3**), MeOH (**PMS2**) and NH₃ (**PAS1**) complexes. The PHA-MeNH₂ complex (**PMAS2**), however, is an exception and once again happens to be a higher energy local minimum.

The difference in stabilization energies for the three isomers of H₂O complexes is about 0.9 kJ mol⁻¹, while in the case of MeOH complexes this difference is only about 0.5 kJ mol⁻¹. However, in the case of NH₃ complexes the difference in the stabilization energy between the global minimum and the next higher energy local minimum is 2.3 kJ mol⁻¹. On the other hand, the observed PHA-MeNH₂ complex (**PMAS2**) is about 4.9 kJ mol⁻¹ higher in energy than global minimum. It must be pointed out here that even though the calculated energy differences between the various isomers is only marginally in the case H₂O and MeOH complexes, while in the case of NH₃ and MeNH₂ complexes the differences are considerable, only one complex was observed experimentally in each case.³⁻⁵

The DFT-SAPT interaction energy decomposition was carried out for the all the structures shown in Figure 2. Table 2 lists the various components of the total energy and also the ZPE corrected SAPT stabilization energies. The ZPE corrected DFT-SAPT stabilization energies are comparable to the ZPE and BSSE corrected CCSD(T) energies

using the same basis set (see Table 1). However, since the energy differences between the various minima are small, the change in the level of calculation leads to rearrangement of relative energies of various isomers in each set. In the case of DFT-SATP calculations the observed complexes of H₂O and NH₃ are the global minima, while the complexes of MeOH and MeNH₂ are higher energy local minima. For the PHA-H₂O system, the observed complex **PWS3** has highest contribution in all the columns, which implies that this structure maximizes all the possible interactions. In the case of MeOH complexes, the induction contribution is almost constant for all the three isomers. The formation of observed **PMS2** structure is favored by the dispersion contribution, while the electrostatic contribution is the lowest among the three isomers. In the case of NH₃ complexes higher contributions from electrostatic and induction energies play dominant role in stabilizing the observed **PAS1** structure. On the other hand, dispersion energy has higher contribution in stabilizing the observed MeNH₂ complex, **PMAS2**. A careful inspection of Table 2 reveals that electrostatics plays major role in stabilizing the observed structures of H₂O and NH₃ complexes, while dispersion plays a major role in stabilizing the observed structures of MeOH and MeNH₂. This implies that the substitution by methyl group switches the lead contribution from electrostatics to dispersion. This, in all probability, can be attributed to the higher polarizability of the O–H and N–H groups in MeOH and MeNH₂, respectively, which can be ascribed to the electron donating ability of the methyl group.

Yet another parameter that was considered was the Gibbs free energy (G). This is due to the fact that the temperature of the experiment is non-zero, therefore the contribution of entropy cannot be neglected. The free energy surface includes the entropy

contribution. The ΔG values were obtained by the thermochemical analysis following vibrational frequency calculations in Gaussian-03 at MP2/aug-cc-pVDZ level of theory and the electronic energy obtained at CCSD(T)/aug-cc-pVDZ level. The Gibbs free energy (ΔG) for the formation of all the complexes was calculated as a function of temperature (T) and the results are presented in Figure 3. Further, the ΔG values at 10 K are listed in Table 1. These plots reveal that very low temperatures (below ~ 100 K) are required for the formation and stability of all the complexes. For the H_2O complexes, below 80 K the ΔG for the formation of **PWS3** is more negative than the other two complexes, and at 10 K, the experimentally observed **PWS3** structure is clearly the most stable complex. In the case of MeOH complexes, the ΔG for the formation of the observed **PMS2** structure is lower than the other two complexes by 0.4 kJ mol^{-1} , which are isoenergetic. With the energy separation of 0.4 kJ mol^{-1} the combined population of the other two species relative to **PMS2** at 10K will be less than 2%. For NH_3 complexes the **PAS1** structure is clearly favored over the other two structures in the entire temperature range. While, in the case of MeNH_2 the observed **PMAS2** is the least favored structure over the entire temperature range. These ΔG calculations clearly establish that at very low temperatures (around 10 K) the observed structures of H_2O (**PWS3**), NH_3 (**PAS1**) and to some extent MeOH (**PMS2**) complexes are thermodynamically the most favored structures. However, it must be noted that in the case of MeOH complex, the difference in the energies of the three is about 0.4 kJ mol^{-1} , which is about the accuracy of the level of calculations reported here. The exclusive observation of **PMS2** structure clearly indicates that there might be effects that clearly cannot be distinguished by the present set of stabilization energy and thermochemical calculations. The exclusive

formation of the **PMS2** structure can perhaps be explained as follows. If we consider a single collision in the molecular beam between PHA and MeOH moieties leads to the formation of the binary complex, then the cross-section for such a collision will be the largest with the π electron density of the benzene ring relative to the two other binding sites. This higher cross-section offered by the π electron density of the benzene ring can be interpreted as an entropic advantage of this site over the other two binding sites. However, it should be noted at this point that the entropy contribution in the thermochemical calculation of ΔG is purely based on the partition function of a particular structure based on rigid rotor – harmonic oscillator – ideal gas approximation (see methodology section), which is different from the entropy based on the interaction cross-section. This entropic advantage in combination with lower ΔG (marginally lower by 0.4 kJ mol⁻¹) manifests in the formation of **PMS2** structure.

Similar arguments can be used in the case of MeNH₂ complex the exclusive formation of **PMAS2** structure can be attributed to the larger cross-section of interaction offered by the π electron density of the benzene ring. However, unlike the MeOH complex (**PMS2**) the MeNH₂ complex (**PMAS2**) is thermodynamically unfavorable over the other two complexes, especially **PMAS1**, which at 10 K is more stable by 4.7 kJ mol⁻¹. This implies that the population of **PMAS2** should be negligibly low. However, the **PMAS2** complex is exclusively formed. Clearly the thermodynamic factors cannot explain such an observation. An alternative explanation based on the kinetics can be as follows: The **PMAS2** structure is a stable minimum and forms during the initial phase of supersonic expansion. However, if the barrier for its inter-conversion to the most stable minimum **PMAS1** is sufficiently high enough to be surmounted at the temperatures

prevailing in the supersonic jet,²⁰ then the PHA-MeNH₂ complex is trapped in the higher energy minimum, which in the present case is **PMAS2**. Therefore, the observation of higher energy **PMAS2** in the present experimental conditions can therefore be attributed to the kinetic trapping.²¹ It must be emphasized here that in the absence of any experimental evidence, this explanation can only at best be tentative.

The ability of formation of H₂O and NH₃ to form the thermodynamically most stable structures can be interpreted on the basis of dominance of electrostatic contribution to the total energy. On the other hand, for the MeOH and MeNH₂ complexes the higher polarizability of the O–H and N–H groups favor interaction with the π electron density of the benzene ring, wherein the dispersion forces dominate. This accompanied by the ability of π electron density of the benzene ring to offer higher cross-section of interaction lead to the formation of MeOH and MeNH₂ complexes. In addition kinetic trapping plays pivotal role in formation of the observed PHA-MeNH₂ complex. Thus in multifunctional molecules the exact hydrogen bonding pattern will be a result of subtle competition between various possibilities, which depends on the fine interplay between the electrostatic and dispersion forces.

Conclusions

The hydrogen-bonded complexes of PHA with H₂O, MeOH, NH₃ and MeNH₂ form a variety of intermolecular structures. The H₂O complex is characterized by the presence of O–H $\cdots\pi$ and C–H \cdots O hydrogen bonds, whilst the O–H group of MeOH interacts with the π electron density of the benzene ring. NH₃ forms a linear C–H \cdots N σ hydrogen-bonded complex. On the other hand MeNH₂ complex is stabilized by the

presence of N–H··· π hydrogen bonding accompanied by a peripheral interaction of methyl C–H group with the acetylenic C \equiv C bond. The stabilization energies calculated at CCSD(T)/aug-cc-pVDZ level along with the calculated values of ΔG for the formation of various complexes reveal that the observed complexes of H₂O and NH₃ are the most stable minima. The MeOH complex is also the most stable minimum; however, its relative stability is only marginally higher than the other two minima. The observed MeNH₂ complex is the highest energy structure both in terms of stabilization and free energies, and definitely is a kinetically trapped structure. The DFT-SAPT calculations indicate that electrostatics play dominant role in stabilizing the observed H₂O and NH₃ complexes, while dispersion is the major contribution for the MeOH and MeNH₂ complexes. The subtle balance between the electrostatic and dispersion forces along with factors such as surface area of interaction and kinetic trapping lead to the observed methyl group induced hydrogen bond switching in the phenylacetylene complexes.

Acknowledgements. Authors wish to thank Mr. Prashant Chandra Singh for the experiments and initial calculations. This material is based upon work supported by Department of Science and Technology (Grant No.SR/S1/PC/23/2008) to G.N.P. This work is part of the research project Z4 055 0506 and is supported by grants from Ministry of Education of the Czech Republic (Center for Biomolecules and Complex Molecular Systems, LC512 and MSM6198959216). The support from Praemium Academiae, Academy of Sciences of the Czech Republic dedicated to P.H. in 2007 is also acknowledged.

References and Notes

- (1) (a) Etter, M. C. *Acc. Chem. Res.* **1990**, 23, 120. (b) Etter, M. C. *J. Phys. Chem.* **1991**, 95, 4601.
- (2) Legon, A. C.; Millen, D. J. *Chem. Soc. Rev.* **1987**, 16, 467.
- (3) Singh, P. C.; Bandyopadhyay, B.; Patwari, G. N. *J. Phys. Chem. A* **2008**, 112, 3360.
- (4) Singh, P. C.; Patwari, G. N. *J. Phys. Chem. A* **2008**, 112, 5121.
- (5) Singh, P. C.; Patwari, G. N. *J. Phys. Chem. A* **2008**, 112, 4426.
- (6) (a) Engdahl, A.; Nelander, B. *J. Phys. Chem.* **1985**, 89, 2860. (b) Wanna, J.; Menapace, J. A.; Bernstein, E. R. *J. Chem. Phys.* **1986**, 85, 1795. (c) Suzuki, S.; Green, P. G.; Bumgarner, R. E.; Dasgupta, S.; Goddard, W. A., III; Blake, G. A. *Science* **1992**, 257, 942. (d) Pribble, R. N.; Garrett, A. W.; Haber, K.; Zwier, T. S. *J. Chem. Phys.* **1995**, 103, 531. (e) Gutowsky, H. S.; Emilsson, T.; Arunan, E. *J. Chem. Phys.* **1993**, 99, 4883.
- (7) (a) Engdahl, A.; Nelander, B. *Chem. Phys. Lett.* **1983**, 100, 129. (b) Peterson, K. I.; Klemperer, W. J. *J. Chem. Phys.* **1984**, 81, 3842.
- (8) Fraser, G. T.; Leopold, K. R.; Klemperer, W. J. *J. Chem. Phys.* **1984**, 80, 1423.
- (9) (a) Jeziorski, B.; Moszynski, R.; Szalewicz, K. *Chem. Rev.* **1994**, 94, 1887. (b) Hesselman, A.; Janssen, G.; Schutz, M. *J. Chem. Phys.* **2005**, 122, 14103.
- (10) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.;

- Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. *Gaussian 03*, Revision A.1; Gaussian, Inc.: Wallingford, CT, 2003.
- (11)MOLPRO, a package of *ab initio* programs designed by Werner H. J. and Knowles, P. J. version 2006.1, Amos, R. D.; Bernhardsson, A.; Berning, A.; Celani, P.; Cooper, D. L.; Deegan, M. J. O.; Dobbyn, A. J.; Eckert, F.; Hampel, C.; Hetzer, G.; Knowles, P. J.; Korona, T.; Lindh, R.; Lloyd, A. W.; McNicholas, S. J.; Manby, F. R.; Meyer, W.; Mura, M. E.; Nicklass, A.; Palmieri, P.; Pitzer, R.; Rauhut, G.; Schu"tz, M.; Schumann, U.; Stoll, H.; Stone, A. J.; Tarroni, R.; Thorsteinsson, T. and Werner, H.-J.
- (12)Stearns, J. A.; Zwier, T. S. *J. Phys. Chem. A*, **2003**, *107*, 10717.
- (13)Singh, P. C.; Patwari, G. N. *Curr. Sci.* **2008**, *95*, 469.
- (14)(a) Siglow, K.; Neusser, H. J. *Chem. Phys. Lett.* **2001**, *343*, 475. (b) Dreizler, H.; Hartke, B.; Rudolph, H. D. *J. Mol. Struct.* **2006**, *825*, 1.

- (15)Lakshmi, B.; Samuelson, A. G.; Jose, K. V. J.; Gadre, S. R.; Arunan, E. *New J. Chem.* **2005**, 29, 371.
- (16)Rodham, D. A.; Suzuki, S.; Suenram, R. D.; Lovas, F. J.; Dasgupta, S.; Goddard, W. A., III; Blake, G. A. *Nature* **1993**, 362, 735.
- (17)The other spectroscopic inputs include the shift in the electronic transitions for the complexes relative to the bare PHA and the IR spectra in the O –H stretching region for the H₂O and MeOH complexes. See references 3-5 for details.
- (18)Dabkowska, I.; Jurecka, P.; Hobza, P. *J. Chem. Phys.* **2005**, 122, 204322.
- (19)(a) Janowski, T.; Pulay, P. *Chem. Phys. Lett.* **2007**, 447, 27. (b) Jurečka, P.; Šponer, J.; Černý, J.; Hobza, P. *Phys. Chem. Chem. Phys.* **2006**, 8, 1985.
- (20)It is difficult to precisely comment on the temperature during the expansion process, but it should lie between the nozzle temperature (298 K) and the terminal temperature (~ 3.5 K for phenylacetylene).
- (21)(a) Douberly, G. E.; Ricks, A. M.; Schleyer, P v. R.; Duncan M. A. *J. Chem. Phys.* **2008**, 128, 021102. (b) Maity, S.; Patwari, G. N. *J. Phys. Chem. A* **2009**, in press.

TABLE 1: ZPE and BSSE Corrected Stabilization Energies (kJ mol⁻¹) of Various PHA Complexes Calculated using aug-cc-pVDZ Basis Set and ΔG for the Formation of Various Complexes at 10K.

	MP2	CCSD(T)	ΔG
PWS1	-6.4	-6.2	-5.6
PWS2	-8.1	-6.5	-6.3
PWS3	-7.6	-7.1	-7.0
PMS1	-9.8	-9.1	-8.6
PMS2	-13.6	-9.6	-8.3
PMS3	-13.1	-9.5	-8.7
PAS1	-8.7	-8.2	-8.3
PAS2	-5.4	-3.2	-2.8
PAS3	-6.7	-5.9	-5.6
PMAS1	-12.4	-11.3	-10.3
PMAS2	-10.6	-6.4	-5.6
PMAS3	-12.1	-9.4	-8.7

TABLE 2: DFT–SAPT Interaction Energy Decomposition (kJ mol⁻¹) for Various Complexes of PHA Calculated using aug-cc-pVDZ Basis Set.

	E _{elec}	E _{ind}	E _{disp}	E _{exch}	δ _{HF}	E _{int}	E _{int} + ZPE
PWS1	-17.0	-2.6	-6.4	17.4	-2.0	-10.6	-6.3
PWS2	-11.9	-2.7	-13.1	18.2	-1.5	-11.0	-7.3
PWS3	-25.5	-4.8	-13.6	32.2	-3.5	-15.2	-9.0
PMS1	-20.9	-3.1	-10.2	24.8	-2.8	-12.2	-9.0
PMS2	-17.4	-3.2	-26.1	36.2	-3.0	-13.5	-10.2
PMS3	-20.2	-3.2	-25.6	37.7	-3.2	-14.6	-10.7
PAS1	-24.9	-4.1	-8.4	27.0	-3.8	-14.2	-8.8
PAS2	-9.3	-1.4	-14.4	19.1	-1.4	-7.4	-3.9
PAS3	-22.4	-3.5	-13.8	29.8	-2.9	-12.8	-7.8
PMAS1	-29.2	-4.7	-12.7	35.9	-5.2	-16.0	-11.8
PMAS2	-15.4	-1.5	-26.3	35.2	-2.6	-10.6	-7.1
PMAS3	-25.5	-3.3	-23.1	41.1	-3.6	-14.4	-10.8




Figure 1. FDIR spectrum of (A) PHA, (B) PHA-Ar, (C) PHA-H₂O, (D) PHA-MeOH, (E) PHA-NH₃ and (F) PHA-MeNH₂, in the acetylenic C–H stretching region. In (A) the arrow indicates the position of the unperturbed C–H oscillator of PHA evaluated using two state deperturbation model (Ref. 13).




Figure 2. Calculated structures of PHA complexes with H₂O, MeOH, NH₃ and MeNH₂ at the MP2/aug-cc-pVDZ level. The distances are shown in Å.




Figure 3. Plot of variation of ΔG vs. T for the formation of various complexes of PHA with (A) H_2O , (B) MeOH , (C) NH_3 and (D) MeNH_2 (see text for details). In each case the solid squares (■) represent the experimentally observed structure.