

ERRATA

- 1) str. 8 – u citace OLIVER & NELSON chybí ročník 1967
- 2) str. 10 – Amblypygi má být bez podtržení
- 3) str. 12 – místo Obr. X má být Obr. 2
- 4) str. 16 – místo citace SUZUKI 1945 má být SUZUKI 1954
- 5) str. 18 – místo viz obr. 6 má být viz obr. 5
- 6) str. 40 – místo viz obr. 11 má být viz. obr. 13; místo viz obr. 12 má být viz obr. 14; místo viz obr. 13 má být viz obr. 15
- 7) str. 55 – věta “ Metodou C-pruhování byl odhalen konstitutivní heterochromatin, který se většinou nacházel...” má být dokončena „ v oblasti centromer.“
- 8) str. 72 – v poslední větě má být odkaz na obr. 48 – viz obr. 48
- 9) str. 75 - místo southernova hybridizace má být Southernova hybridizace
- 10) kompletní seznam literatury:

Ahn, I.Y., Winter, C.E. (2006): The genome of *Oscheius Tipulae*: Determination of size, complexity, and structure by DNA reassociation using fluorescent dye. *Genome* 49:1007-1015.

Angelier, N., Tramier, M., Louvet, E., Coppey-Moisan, M., Savino, T. M., De Mey, J. R., Hernandez-Verdun, D. (2005): Tracking the interactions of rRNA processing proteins during nucleolar assembly in living cells. *Molecular Biology of the Cell* 16:2862-2871.

Araújo, D., Brescovit, A. D., Rheims, C. A. & Cella, D. M. (2005): Chromosomal data of two pholcids (Araneae, Pholcidae): A new diploid number and the first cytogenetical record for the New World clade. *Journal of Arachnology* 33:591-596.

Araújo, D., Rheims, C.A., Brescovit, A.D., Cella, D.M. (2008) Extreme degree of chromosome number variability in species of the spider genus *Scytodes* (Araneae, Haplogynae, Scytodidae). *Journal of Zoological Systematics and Evolutionary Research* 46:89-95.

Ax, P. (1996): *Multicellular animals. A new approach to the phylogenetic order in nature.* Springer, Berlin, Heidelberg, New York:1-225.

Azzalin, C.M., Nergadze, S.G., Giulotto, E. (2001): Human intrachromosomal telomeric-like repeats: sequence organization and mechanisms of origin, *Chromosoma* 110:75–82.

Bachtrog, D. (2008): The temporal dynamics of processes underlying Y chromosome degeneration. *Genetics* 179:1513-1525.

Bailey, S.M., Goodwin, E.H., Cornforth, M.N. (2004): Strand-specific fluorescence in situ hybridization: the CO-FISH family. *Cytogenetic and Genome Research* 107:14-17.

Baird, D.M. (2005): New developments in telomere length analysis. *Experimental gerontology* 40:363-368.

- Balajee, A.S., Natarajan, H.J. Oh, A.T. (1994): Analysis of restriction enzyme-induced chromosome aberrations in the interstitial telomeric repeat sequences of CHO and CHE cells by FISH, *Mutation Research* 307:307–313.
- Benavente, R., Wettstein, R. (1980): Ultrastructural characterization of the sex chromosomes during spermatogenesis of spiders having holocentric chromosomes and a long diffuse stage. *Chromosoma* 77: 69-81.
- Bertoni, L., Attolini, C., Tessera, L., Mucciolo, E. Giulotto, E (1994): Telomeric and nontelomeric (TTAGGG)_n sequences in gene amplification and chromosome stability. *Genomics* 24:53-62.
- Bi, K., Bogart, J.P. (2006): Identification of intergenomic recombinations in unisexual salamanders of the genus *Amphispiza* by genomic in situ hybridization (GISH). *Cytogenetic and Genome Research* 112: 307-312.
- Biessmann, H., Mason, J.M. (2003): Telomerase-independent mechanisms of telomere maintenance. *Cellular and Molecular Life Sciences* 60:2325-2333.
- Bole-Gowda, B.N. (1952): Studies on the chromosomal and sex – determining mechanism of four hunting spiders (Sparassidae). *Proceedings of Zoological Society of Bengal* 5:51-70.
- Bolzán, A.D., Bianchi, M.S. (2006): Telomeres, interstitial telomeric repeat sequences, and chromosomal aberrations. *Mutation Research* 612:189-214.
- Brown, D. D., and Gurdon, J. B. (1964): Absence of ribosomal RNA synthesis in the anucleolate mutant of *Xenopus Laevis*. *Proceeding of the National Academy of Science of the USA* 51:139-146.
- Bull, J.J. (1983): *Evolution of Sex Determining Mechanisms*. Benjamin/Cummings, Menlo Park, CA, USA.
- Cabrero, J., Camacho, J.P.M. (2008): Location and expression of ribosomal RNA genes in grasshoppers: abundance of silent and cryptic loci. *Chromosome Research* 16:595-607.
- Carvalho, B.A., Koerich, L.B., Clark, A.G. (2009): Origin and evolution of Y chromosomes: *Drosophila* tales. *Trends in Genetics* 25:433-440.
- Castro, L.F.C., Holland, P.W.H. (2002): Fluorescent in situ hybridisation to amphioxus chromosomes. *Zoological Science* 19:1349-1353.
- Cesare, A.J., Quinney, N., Willcox, S., Subramanian, D., Griffith, J.D. (2003): Telomeres looping in *P. sativum* (Common garden pea). *Plant Journal* 36:271-79.
- Ciufo, L.F., Brown, J.D. (2000): Nuclear export of yeast signal recognition particle lacking Srp54p by the Xpo1p/Crm1p NES-dependent pathway. *Current Biology* 10:1256-1264.
- Coddington, J.A., Levi, H.W. (1991): Systematic and evolution of spiders (Araneae): *Annual Review of Ecology and Systematics* 22:565-592.
- Coghill, G., Grant, A., Orrell, J.M., Jankowski, J. Evans, A.T. (1990): Improved silver staining of nucleolar organiser regions in paraffin wax sections using an inverted incubation technique. *Journal of Clinical Pathology* 43:1029-1031.

Cox, A.V., Bennett, S.T., Parokony, A.S., Kenton, A., Callimassia, M.A., Bennett, M.D. (1993): Comparison of plant telomere locations using a PCR-generated synthetic probe. *Annals of Botany* 72:239-247.

Datta, S.N., Chatterjee, K. (1988): Chromosomes and sex determination in 13 araneid spiders of North-Eastern India. *Genetica* 76: 91-99.

Dechampsme, A. M., Koroleva, O., Leger-Silvestre, I., Gas, N., and Camier, S. (1999): Assembly of 5S ribosomal RNA is required at a specific step of the pre-rRNA processing pathway. *Journal of Cell Biology* 145:1369-1380.

de Sá, R.O., Hillis, D.M. (1990): Phylogenetic relationship of the pipid frogs *Xenopus* and *Silurana*: an integration of ribosomal DNA and morphology. *Molecular Biology and Evolution* 7:365-376.

DiMario, P. J. (2004): Cell and molecular biology of nucleolar assembly and disassembly. *International Review of Cytology* 239:99-178.

Dolejš, P., Kořínková, T., Musilová, J., Opatová, V., Kubcová, L., Buchar, J., Král, J. (2011): Karyotypes of central European spiders of the genera *Arctosa*, *Tricca*, and *Xerolycosa* (*Araneae*: *Lycosidae*). *European Journal of Entomology* 108:1-16.

Dulíková, L. & Král, J. (2007): Insights into the karyotype evolution of the spider infraorder Mygalomorphae. In: Pinto-da-Rocha R., Machado. G., Brescovit, A.D.(eds): Abstracts, 17th International Congress of Arachnology, Sao Paulo.

Dundr, M., Olson, M.O.J. (1998): Partially processed pre-rRNA is preserved in association with processing components in nucleolus-derived foci during mitosis. *Molecular Biology of the Cell* 9:2407-2422.

Dunlop, J.A. (1996): Evidence for a sister group relationship between Ricinulei and Trigonotarvida. *Bulletin of the British Arachnological Society* 10:193-204.

Eisner, T., Meinwald, J., Monro, A. Ghent, R. (1961): Defense mechanisms of arthropods - I. The composition and function of the spray of the whipscorpion, *Mastigoproctus giganteus* (Lucas) (Arachnida: Pedipalpida). *Journal of Insect Physiology* 6:272-298.

Eliceiri, G.L. (1999): Small nucleolar RNAs. *Cellular and Molecular Life Sciences* 56:22-31.

Ellegren, H. (2011): Sex-chromosome evolution: recent progress and the influence of male and female heterogamety. *Genetics* 12:157-166.

Fahrein, K., Masta, S.E., Podsiadlowski, L. (2009): The first complete mitochondrial genome sequences of *Amblypygi* (Chelicerata: Arachnida) reveal conservation of the ancestral arthropod gene order. *Genome* 52: 456-466.

Fajkus, J., Sykorova, E., Leitch, A.R. (2005): Technique in plant telomere biology. *Biotechnologies* 38: 233-243.

Fang, G., Cech, T. R. (1993a): The β subunit of oxytricha telomere-binding protein promotes G-quartet formation by telomeric DNA. *Cell* 74:875-885.

Fang, G., Cech, T. R. (1993b): Oxytricha telomere-binding protein: DNA-dependent dimerization of the α and β subunits. *Proceedings of the National Academy of Sciences of the USA* 90:6056-6060.

- Fatica, A., Tollervey, D. (2002): Making ribosomes. *Current Opinion in Cell Biology* 14:313-318.
- Felsenstein, J. (1974): The evolutionary advantage of recombination. *Genetics* 78:737-756.
- Forman, M., Nguyen, P., Hula, V., Král, J. (2013): Sex chromosome pairing and extensive NOR polymorphism in *Wadicosa fidelis* (Araneae: Lycosidae). *Cytogenetic Geome Research* 141:43-49.
- Fuchs, J., Brandes, A., Schubert, I. (1995): Telomere sequence localization and karyotype evolution in higher plants. *Plant Systematics and Evolution* 196:227-241.
- Furlong, J.C., Maden, B.E.H. (1983): Patterns of major divergence between the internal transcribed spacers of ribosomal DNA in *Xenopus borealis* and *Xenopus laevis*, and of minimal divergence within ribosomal coding regions. *EMBO Journal* 2:443-448.
- Gautier, T., Robert-Nicoud, M., Guilly, M., Hernandez-Verdun, D. (1992): Relocation of nucleolar proteins around chromosomes at mitosis: A study by confocal laser scanning microscopy. *Journal of Cell Science* 102: 729-737.
- Gerbi, S.A., and Borovjagin, A.V. (2004): Pre-ribosomal RNA processing in multicellular organisms. In "The Nucleolus" (M.O. J. Olson, Ed.): 170-198.
- Glover, T.W., Stein, C.K. (1988): Chromosome breakage and recombination at fragile sites. *American Journal of Human Genetic* 43:265-273.
- Goodwin, E., Meyne, J. (1993): Strand-specific fish reveals orientation of chromosome-18 aliphoid dna. *Cytogenetics and Cell Genetics* 63:126-127.
- Graham, D.E. (1978): The isolation of high molecular weight DNA from whole organisms or large tissue masses. *Analytical Biochemistry* 85:609-613.
- Greider, C.W., Blackburn, E.H. (1989): A telomeric sequence in the RNA of tetrahymena telomerase required for telomere repeat synthesis. *Nature* 337:331-337.
- Griffith, J.D., Comeau, L., Rosenfield, S., Stansel, R.M., Bianchi, A., Moss, H., De Lange, T. (1999): Mammalian telomeres end in a large duplex loop. *Cell* 97:503-514.
- Griffiths, A.P., Butler, C.W., Roberts, P., Dixon, M.F., Quirke, P. (1989): Silver stained structures (AgNORs) their dependence on tissue fixation and absence of prognostic relevance in rectal adenocarcinoma. *Journal of Pathology* 159:121-127.
- Gutell, R.R., Fox, G.E. (1988): A compilation of large subunit RNA sequences presented in a structural format. *Nucleic Acids Research* 16:75-269.
- Hackman, W. (1948): Chromosomenstudien an Araneen mit besonderer Berücksichtigung der Geschlechtschromosomen. *Acta Zoologica Fenica* 54:1-101.
- Hadjiolov, A.A. (1985): „The nucleolus and ribosome biogenesis“ Springer, New York.
- Halanych, K.M. (1991): 5S ribosomal RNA sequences inappropriate for phylogenetic reconstruction. *Molecular Biology and Evolution* 8:249-253.
- Harvey, M.S. (2002): The neglected cousins: what do we know about the smaller arachnid orders? *Journal of Arachnology* 30:357-372.

Harvey, M.S. (2003): Catalogue of the smaller arachnid orders of the World: Amblypygi, Uropygi, Schizomida, Palpigradi, Ricinulei and Solifugae. CSIRO Publishing, Collingwood, Victoria, Australia.

Harvey, M.S. (2007): The smaller arachnid orders: diversity, descriptions and distributions from Linnaeus to the present (1758 to 2007). *Zootaxa* 1668:363-380.

Haupt, J., Höhne, G., Schwarz, H., Chen, B., Zhao, W., Zhang, Y. (1988): Chinese whip scorpions using 2-ketones in defense secretion (Arachnida:Uropygi). *Journal of Comparative Physiology B* 157: 883-885.

Haupt, J., Höhne, G., Weiske, T. (1993): Acetic acid esters, *n*-hexanol, *n*-octanol, and capronic acid as ingredients in the defense secretion product of whip scorpions. C. R. XIVE Coll. Europ. Arachnol. Catania, 23D27 aout 1993. *Bollettino dell'Accademia Gioenia Scienze Naturali* 26:175-180.

Heitz, E. (1931): Die ursache der gesetzmässigen zahl. lage, form und grösse pflanzlicher nukleolen. *Planta* 12:775-884.

Hernandez-Verdun, D., Roussel, P., Gautier, T. (1993): Nucleolar proteins during mitosis. *Chromosomes Today* 11:79-90.

Hill, C.A., Guerrero, F.D., Van Zee, J.P., Geraci, N., Walling, J.G., Stuart, J.J. (2009): The position of repetitive DNA sequence in the southern cattle tick genome permits chromosome identification. *Chromosome Research* 17:77-89.

Hillis, D.M., Dixon, M.T. (1989): Vertebrate phylogeny: evidence from 28S ribosomal DNA sequences. In B. Fernholm, K. Bremer, and H. Jornvall (eds.), *The Hierarchy of Life*:355-367.

Hillis, D.M., Dixon M.T. (1991): Ribosomal DNA: molecular evolution and phylogenetic inference. *Quarterly Review of Biology* 66:411-453.

Hillis, D.M., Dixon, M.T., Ammerman, L.K. (1991): The relationships of the coelacanth *Latimeria chalumnae*: evidence from sequences of vertebrate 28S ribosomal RNA genes. *Environmental Biology of Fishes* 32:119-130.

Hillis, D.M., Larson, A., Davis, S.K., Zimmer, E.A. (1990): Nucleic Acids III: sequencing. In D. M. Hillis and C. Moritz (eds.), *Molecular Systematics*:318-370.

Hirai, H., LoVerde, P.T. (1996): Identification of the telomeres on *Schistosoma mansoni* chromosomes by FISH. *Journal of Parasitology* 82:511-512.

Hiscox, J.A. (2002): Brief review: The nucleolus—a gateway to viral infection? *Archives of Virology* 147:1077-1089.

Hofgärtner, F.J., Schmid, M., Krone, W., Zenzes, M.T., Engel, W. (1979): Pattern of activity of nucleolus organizers during spermatogenesis in mammals as analyzed by silver-staining. *Chromosoma* 71:197-216.

Hoshikawa, Y., Iida, Y., Iwabuchi, M. (1983): Nucleotide sequence of the transcriptional initiation region of *Dictyostelium discoideum* rRNA gene and comparison of the initiation regions of three lower eukaryotes' genes. *Nucleic Acids Research* 11:1725-1734.

Howell, W.M. & Black, D.A. (1980): Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36: 1014.

Charlesworth, B. (1996): The evolution of chromosomal sex determination and dosage compensation. *Current Biology* 6: 149-162.

Charlesworth, B., Charlesworth, D. (2000): The degeneration of Y chromosomes. *Philosophical Transactions of the Royal Society B: Biological Sciences* 355:1563-1572.

Charlesworth, D., Charlesworth, B. (2005): Sex chromosomes: Evolution of the weird and wonderful. *Current Biology* 15:129-131.

Chen, D., Dundr, M., Wang, C., Leung, A., Lamond, A., Misteli, T., Huang, S. (2005): Condensed mitotic chromatin is accessible to transcription factors and chromatin structural proteins. *Journal of Cell Biology* 168:41-54.

Jablonka, E., Lamb, M.J. (1988): Meiotic pairing constraints and the activity of sex chromosomes. *Journal of Theoretical Biology* 133:23-36.

Jaffe, K. (2004): Sex promotes gamete selection: A quantitative comparative study of features favoring the evolution of sex. *Complexity* 9:43-51.

Jha, A.N., Dominiquez, I., Balajee, A.S., Hutchinson, T.H., Dixon, D.R., Natarajan, A.T. (1995): Localization of a vertebrate telomeric sequence in the chromosomes of two marine worms. (phylum Annelida, class Polychaeta). *Chromosome Res* 3: 507-508.

Joffe, B.I., Solovei, I.V., Macgregor, H.C. (1998): Ends of chromosomes in *Polycelis tenuis* (Platyhelminthes) have telomere repeat TTAGGG. *Chromosome Research* 4:323-324.

Jiménez-García, L.F., Segura-Valdez, M.D.L., Ochs, R.L., Rothblum, L.I., Hannan, R., Spector, D.L. (1994): Nucleologenesis: U3 snRNA-containing prenucleolar bodies move to sites of active pre-rRNA transcription after mitosis. *Molecular Biology of the Cell* 5:955-966.

Kahn J., Quart, J. (1964): Cytotaxonomy of ticks. *Quarterly Journal of Microscopical Science* 105:123-137.

Kasturi Bai, A.R. & Parthasarathy, M.D. (1957): The chromosomes of *Thelyphonus indicus* Stoliczka. – *Proceeding of the National Institute of Science of India - Section B* 24:19-22.

Khurts, S., Masutomi, K., Delgermaa, L., Arai, K., Oishi, N., Mizuno, H., Hayashi, N., Hahn, W.C., Murakami, S. (2004): Nucleolin interacts with telomerase. *Journal of Biological Chemistry* 279:51508-15.

Kilburn, A.E., Shea, M.J., Sargent, R.G., Wilson, J.H. (2001): Insertion of a telomere repeat sequence into a mammalian gene causes chromosome instability. *Molecular and Cellular Biology* 21:126–135.

Kilian, A., Kudrna, D., Kleinhofs, A. (1999): Genetic and molecular characterization of barley chromosome telomeres. *Genome* 42:412–419.

Kiss, T. (2002): Small nucleolar RNAs: An abundant group of noncoding RNAs with diverse cellular functions. *Cell* 109:145–148.

Koch, A.M., Reynolds, F., Kircher, M.F., Merkle, H.P., Weidleder, R., Josephson, L. (2003): Uptake and metabolism of dual fluorochrome tat-nanoparticle in HeLa cells. *Bioconjugate Chemistry* 14: 1115-1121.

Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Paabo, S., Villablanca, F.X., Wilson, A.C. (1989): Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences of the USA* 86:6196-6200.

Korek, G. Martin, H., Wenzelides, K. (1991): A modified method for the detection of nucleolar organizer regions (AgNORs). *Acta Histochemica* 90:155-157.

Král, J. (2007): Evolution of multiple sex chromosomes in the spider genus *Malthonica* (Araneae: Agelenidae) indicates unique structure of the spider sex chromosome systems. *Chromosome Research* 15: 863-879.

Král, J., Musilová, J., Šťáhlavský, F., Řezáč, M., Akan, Z., Edwards, R.L., Coyle, F.A., Ribera Almeje, C. (2006): Evolution of the karyotype and sex chromosome systems in basal clades of araneomorph spiders (Araneae: Araneomorphae). *Chromosome Research* 14: 859-880.

Král, J., Kováč, L., Šťáhlavský, F., Lonský, P., Luptáček, P. (2008): The first karyotype study in palpigrades, a primitive order of arachnids (Arachnida: Palpigradi). *Genetica* 134:79-87.

Král J., Kořínková T., Forman M., Krkavcová L. (2011): Insights into the meiotic behavior and evolution of multiple sex chromosome systems in spiders. *Cytogenetic and Genome Research* 133: 43-66.

Král, J., Kořínková, T., Krkavcová, L., Musilová, J., Forman, M., Ávila, I., Haddad, C.R., Vítková, M., Henriques, S., Palacios, J., Hedin, M. (2013): Evolution of karyotype, sex chromosomes, and meiosis in mygalomorph spiders (Araneae: Mygalomorphae). *Biological Journal of the Linnean Society* 109:377-408.

Kubai, D.F., Wise, D. (1981): Nonrandom chromosome segregation in *Neocurtilla* (*Gryllotalpa*) *hexadactyla*: An ultrastructural study. *Journal of Cell Biology* 88:281-293.

Laird, D.J., Weissman, I.L. (2004): Telomerase maintained in selfrenewing tissues during serial regeneration of the urochordate *Bothryllus schosserii*. *Developmental Biology* 273:185-194.

Larson, A., Wilson, A.C. (1989): Patterns of ribosomal RNA evolution in salamanders. *Molecular Biology and Evolution* 6:131-154.

Leung, A.K.L., Gerlich, D., Miller, G., Lyon, C., Lam, Y.W., Lleres, D., Daigle, N., Zomerdijs, J., Ellenberg, J., Lamond, A.I. (2004): Quantitative kinetic analysis of nucleolar breakdown and reassembly during mitosis in live human cells. *Journal of Cell Biology* 166:787-800.

Leung, A.K.L., Lamond, A.I. (2003): The dynamics of the nucleolus. *Critical Reviews in Eukaryotic Gene Expression* 13:39-54.

Lebovsky, R., Bensimon, A. (2005): DNA replication origin plasticity and perturbed fork progression in human inverted repeats. *Molecular Cell Biology* 25: 6789-6797.

Levan, A.K., Fredga K., Sandberg A.A. (1964): Nomenclature for centromeric position on chromosomes. *Hereditas* 52:201-220.

Lin, K.W., Yan, J. (2005): The telomere length dynamic and methods of its assessment. *Journal of Cellular and Molecular Medicine* 9:977-989.

Lin, K.W., Yan, J (2008): Endings in the middle: Current knowledge of interstitial telomeric sequences. *Mutation Research* 658:95-110.

- Lockwood, A.M.P. (1961): „Ringer“ solutions and some notes on the physiological basis their ionic composition. *Comparative Biochemistry and Physiology* 2:241-289.
- Long, E.O., Dawid, I.B. (1980): Repeated genes in eukaryotes. *Annual Review of Biochemistry* 49: 727-764.
- Lu, W., Zhang, Y., Liu, D., Songyang, Z., Wan, M. (2013): Telomeres-structure, function, and regulation. *Experimental Cell Research* 319:133-141.
- McClintock, B. (1934): The relationship of particular chromosomal element to the development of the nucleoli in *Zea Mays*. *Zeitschrift für Zellforschung und Mikroskopische Anatomie* 21:294-328.
- McClintock, B. (1941): The stability of broken ends of chromosomes in *Zea Mays*. *Genetics* 26:234-282.
- McLemore, D.D., Lord, B.A. (1991): Argyrophilic nucleolar organizer regions - an improved technique for visualization. *Journal of Histotechnology* 14:187-189.
- Mallatt, J., Craig, C.W., Yoder, M.J. (2010): Nearly complete rRNA genes assembled from across the metazoan animals: Effects of more taxa, a structure-based alignment, and paired-sites evolutionary models on phylogeny reconstruction. *Molecular Phylogenetics and Evolution* 55: 1-17.
- Mason, J. M., Biessmann, H. (1995): The unusual telomeres of drosophila. *Trends in Genetics* 11:58-62.
- Meyne, J., Baker, R.J., Hobart, H.H., Hsu, T.C., Ryder, O.A., Ward, O.G., Wiley, J.E., Wurster-Hill, D.H., Yates, T.L., Moyzis, R.K. (1990): Distribution of non-telomeric sites of the (TTAGGG)_n telomeric sequence in vertebrate chromosomes. *Chromosoma* 99:3-10.
- Meyne, J., Ratliff, R. L., Moyzis, R. K. (1989): Conservation of the human telomere sequence (TTAGGG)_n among vertebrates. *Proceedings of the National Academy of Sciences of the USA* 86:7049-7053.
- Mirre, C., Stahl, A. (1976): Ultrastructural study of nucleolar organizers in the quail oocyte during meiotic prophase. *Journal of Ultrastructure Research* 56:186-201.
- Moyzis, R.K., Buckingham, J.M., Cram, L.S., Dani, M., Deaven, L.L., Jones, M.D., Meyne, J., Ratliff, R.L., Wu, J.R. (1988): A highly conserved repetitive DNA sequence, (TTAGGG)_n, present at the telomeres of human chromosomes. *Proceedings of the National Academy of Sciences of the USA* 85:6622-6626.
- Muller, F., Wicky, C., Spicher, A., Tobler, H. (1991): New telomere formation after developmentally regulated chromosomal breakage during the process of chromatin diminution in *ascaris lumbricoides*. *Cell* 67:815-822.
- Muller, H.J. (1938): The remaking of chromosomes. *Collecting net* 13:181-195.
- Munoz – Jordan, J.L., Cross, J.A., De Lange, T., Griffith, G.D. (2001): T loops at trypanosome and telomeres. *EMBO Journal* 20:579-88.
- Nazar, R.N., Sitz, T.O., Busch, H. (1976): Sequence homologies in mammalian 5.8S ribosomal RNA. *Biochemistry* 15:505-508.

- Niedermaier, J., Moritz, K.B. (2000): Organization and dynamics of satellite and telomere DNAs in ascaris: Implications for formation and programmed breakdown of compound chromosomes. *Chromosoma* 109:439-452.
- Nikicicz, E.P., Norback, D.H. (1991): Methods in pathology. The use of a colloidal silver (AgNOR) method in assessing bone marrow iron stores and sideroblasts. *Modern Pathology* 4:363-367.
- Norton, R.A., Kethley, J.B., Johnston, D.E. & O'Connor, B.M.(1993): Phylogenetic perspectives on genetic systems and reproductive modes of mites. In: Wrensch D, Ebbert M (eds.) *Evolution and Diversity of Sex Ratio in Insects and Mites*. Chapman & Hall, London.
- Okazaki, S., Tsuchida, K., Maekawa, H., Ishikawa, H., Fujiwara, H. (1993): Identification of a pentanucleotide telomeric sequence, (TTAGG)_n, in the silkworm *bombyx mori* and in other insects. *Molecular and Cellular Biology* 13:1424-1432.
- Oliver, J.H. (1977): Cytogenetics of mites and ticks. *Annual Review of Entomology* 22:407-429.
- Oliver, J.H., Nelson, B.C.(1967): Mite chromosomes: an exceptionally small number. *Nature* 214: 809.
- Olovnikov, A.M. (1973): A theory of marginotomy: The incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *Journal of Theoretical Biology* 41:181-190.
- Olson, M.O., Dundr, M. (2005): The moving parts of the nucleolus. *Histochemistry and Cell Biology*12:203-216.
- Olson, M.O., Hingorani, K., Szebeni, A. (2002): Conventional and nonconventional roles of the nucleolus. *International Review of Cytology* 219:199-266.
- Palm, W., De Lange, T. (2008): How shelterin protects mammalian telomere. *Annual Review of Genetics* 42:301-334.
- Patau K. (1948): X-segregation and heterochromasy in the spider *Aranea reaumuri*. *Heredity* 2: 77-100.
- Pederson, T. (1998): The plurifunctional nucleolus. *Nucleic Acids Research* 26:3871-3876.
- Pellestor, F. and Paulasová, P. (2004): Peptids nucleic acid (PNAs): Introduction to a new class of probes for chromosomal investigation. *Chromosoma* 112: 375-380.
- Petrunkevitch, A. (1955): Arachnida. In Moore, R.C. (Ed.), *Treatise on invertebrate Paleontology*. University of Kansas Press, Lawrence, Kansas.
- Platnick, N.I. (2013): The world spider catalogue, version 14.0. American Museum of Natural History. Přístupno na <http://research.amnh.org/iz/spiders/catalog/>
- Plohl, M., Prats, E., Martinez-Lage, A., Gonzalez-Tizon, A., Mendez, J., Cornudella, L. (2002): Telomeric localization of the vertebrate type hexamer repeat, (TTAGGG)_n, in the wedge shell clam *Donax trunculus* and other marine invertebrate genomes. *Journal of Biological Chemistry* 277:19839-19846.
- Postiglioni, A., Brum-Zorilla, N. (1981): Karyological studies on Uruguayan spiders. II. Sex chromosomes in spiders of the genus *Lycosa* (Araneae:Lycosidae). *Genetica* 56: 47-53.

- Rhodes, D., Giraldo, R. (1995): Telomere structure and function. *Current Opinion in Structural Biology* 5:311-322.
- Rosén, M., Edström, J. (2000): DNA structures common for chironomid telomeres terminating with complex repeats. *Insect Molecular Biology* 9:341-347.
- Rice, W.R. (1996): Evolution of the Y sex chromosome in animals. *Bioscience* 46:331-343.
- Rivero, M.T., Mosquera, A., Goyanes, V., Slijepcevic, P., Fernandez, J.L. (2004): Differences in repair profiles of interstitial telomeric sites between normal and DNA double-strand break repair deficient Chinese hamster cells. *Experimental Cell Research* 295:161-172.
- Rodríguez-Gil, S.G., Merani, M.S., Scioscia, C.L., Mola, L.M. (2007): Cytogenetics in three species of *Polybetes* Simon 1897 from Argentina (Araneae, Sparassidae) I. Karyotype and chromosome banding pattern. *Journal of Arachnology* 35:227-237.
- Rubbi, C.P., Milner, J. (2003): Disruption of the nucleolus mediates stabilization of p53 in response to DNA damage and other stresses. *EMBO Journal* 22:6068-6077.
- Ruiz-Herrera, A., Nergadze, S.G., Santagostino, M., Guilotto, E. (2008): Telomeric repeats far from the ends: mechanisms of origin and role in evolution. *Cytogenetic and Genome Research* 122:219-228.
- Řezáč, M., Král, J., Musilová, J., Pekár, S. (2006): Unusual karyotype diversity in the European spiders of the genus *Atypus* (Araneae: Atypidae). *Hereditas* 143:123-129.
- Sahara, K., Marec, F., Traut, W. (1999): TTAGG telomeric repeats in chromosomes of some insects and other arthropods. *Chromosome Research* 7:449-460.
- Saldanha, S.N., Andrews, L.G., Tollefsbol, T.O. (2003): Assessment of telomere length and factors that contribute to its stability. *European Journal of Biochemistry* 270:389-403.
- Sansam, C.L., Wells, K.S., Emeson, R.B. (2003): Modulation of RNA editing by functional nucleolar sequestration of ADAR2. *Proceedings of the National Academy of Science of the USA* 100:14018-23.
- Sato, S. (1988): Cytological evidence on the ability of the nucleolus organizing regions to assemble pre-existing nucleolar material. *Experientia* 44:264-266.
- Satya-Prakash, K.L., and Pathak, S. (1984): Silver staining pattern of male meiosis in the house cricket. *Journal of Heredity* 75:319-320.
- Savino, T.M., Gébrane-Younès, J., De Mey, J., Sibarita, J., Hernandez-Verdun, D. (2001): Nucleolar assembly of the rRNA processing machinery in living cells. *Journal of Cell Biology* 153:1097-1110.
- Selden, P.A., Shear, W.A., Bonamo, P.M. (1991): A spider and other arachnids from the devonian of New-York, and reinterpretations of devonian araneae. *Palaeontology* 34: 241-281.
- Selden, P.A. (1993) Fossil arachnids-recent advances and future prospects. *Memoirs of the Queensland Museum* 33:389-400.
- Sember, A. (2010): Analýza karyotypu u vybraných bičovců řádů Amblypygi a Uropygi. Diplomová práce. Přírodovědecká fakulta Univerzity Karlovy, Praha.

- Shanahan, C.M. (1989): Cytogenetics of Australian scorpions. II. Chromosome polymorphism in species of *Urodacus* (family Scorpionidae). *Genome* 32:890-900.
- Shear, W.A., Selden, P.A., Rolfe, W.D.I., Bonamo, P.M., Grierson, J.D. (1987): New terrestrial arachnids from the Devonian of Gilboa, New York (Arachnida, Trigonotarbida). *American Museum of Novitates* 2901:1-74.
- Sheldon, S., Lehman, J. M. (1981): Continued presence of nucleoli in human germ cell tumors during mitosis. *Cancer Research*, 41:3666-3668.
- Shultz, J.W. (1989): Morphology of locomotor appendages in Arachnida: evolutionary trends and phylogenetic implications. *Zoological Journal of the Linnean Society (London)* 97:1-56.
- Shultz, J.W. (1990): Evolutionary morphology and phylogeny of Arachnida. *Cladistics* 6:1-38.
- Shultz, J.W. (2007): A phylogenetic analysis of the arachnid orders based on morphological characters. *Zoological Journal of the Linnean Society (London)* 150:221-265.
- Schmid, M., Löser, C., Schmidtke, J., Engel, W. (1982): Evolutionary conservation of a common pattern of activity of nucleolus organizers during spermatogenesis in vertebrates. *Chromosoma* 86:149-179.
- Schneider, M.C., Zacaro, A.A., Oliveira, R.M., Gnaspini, P., Cella, D.M. (2008): Conventional and ultrastructural analyses of the chromosomes of *Discocyrtus pectinifemur* (Opiliones, Laniatores, Gonyleptidae). *Journal of Zoological Systematics and Evolutionary Research* 47:203-207.
- Schneider, M.C., Zacaro, A.A., Pinto-Da-Rocha, R., Candido, D.M., Cella, D.M. (2009): A comparative cytogenetic analysis of 2 Bothriiidae species and overview of the chromosome data of scorpions. *Journal of Heredity* 100:546-555.
- Schneider, M.C., Cella, D.M. (2010): Karyotype Conservation in 2 Populations of the Parthenogenetic Scorpion *Tityus serrulatus* (Buthidae): rDNA and Its Associated Heterochromatin Are Concentrated on Only One Chromosome. *Journal of Heredity* 101:491-496.
- Schubert, I., Rieger, R., Fuchs, J. (1995): Alteration of basic chromosome number by fusion–fission cycles. *Genome* 38:1289-1292.
- Schwander, T., Beukeboom, L.W. (2011): Non-random autosome segregation: A stepping stone for the evolution of sex chromosome complexes?: Sex-biased transmission of autosomes could facilitate the spread of antagonistic alleles, and generate sex-chromosome systems with multiple X or Y chromosomes. *BioEssays* 33:111-114.
- Schwarzacher, H.G., Mikelsaar, A.V., Schnedl, W. (1978): Human nucleolus organizers. *Verhandlungen der Anatomischen Gesellschaft* 72:367-368.
- Schwarzacher, H.G., Wachtler, F. (1983): Nucleolus organizer and nucleoli. *Human Genetics* 63: 89-99
- Simon, C., Franke, A., Martin, A. (1991): The polymerase chain reaction: DNA extraction and amplification. In G. M. Hewitt, A. W. B. Johnston, and J. P. W. Young (eds.). *Molecular Techniques in Taxonomy*: 329-355.
- Sirri, V., Roussel, P., Hernandez-Verdun, D. (2000): The AgNOR proteins: Qualitative and quantitative changes during the cell cycle. *Micron* 31:121-126.

- Slijepcevic, P., Xiao, Y., Dominguez, I., Natarajan, A.T. (1996): Spontaneous and radiation-induced chromosomal breakage at interstitial telomeric sites, *Chromosoma* 104:596-604.
- Sommerville, J., Brumwell, C. L., Politz, J. C., and Pederson, T. (2005): Signal recognition particle assembly in relation to the function of amplified nucleoli of *Xenopus* oocytes. *Journal of Cell Science* 118:1299-1307.
- Soyer-Gobillard, M., Geraud, M. (1992): Nucleolus behaviour during the cell cycle of a primitive dinoflagellate eukaryote, *prorocentrum micans* ehr., seen by light microscopy and electron microscopy. *Journal of Cell Science* 102:475-485.
- Steele, K.P., Holsinger, K.E., Jansen, R.K., Taylor, D.W. (1991): Assessing the reliability of 5S rRNA sequence data for phylogenetic analysis in green plants. *Molecular Biology and Evolution* 8: 240-248.
- Steinemann, S., Steinemann, M. (2005): Retroelements: Tools for sex chromosome evolution. *Cytogenetic and Genome Research* 110:134-143.
- Strunnikov, A. V. (2005): A case of selfish nucleolar segregation. *Cell Cycle* 4:113-117.
- Sturm, H. (1958): Indirekte Spermatophorenubertragung bei dem Geisselskorpion *Trithyreus sturmi* Kraus (Schizomidae, Pedipalpi). *Naturwissenschaften* 45:142-143.
- Sumner, A.T. (1990): Chromosome banding. Unwin hyman, London.
- Sumner, A.T. (2003): Chromosomes: organization and function. Blackwell Science, Oxford
- Suzuki, S. (1954): Cytological studies in spiders III. Studies on the chromosomes of fifty-seven species of spiders belonging to seventeen families, with general considerations on chromosomal evolution. *Journal of Science of the Hiroshima University* 15:23-136.
- Šťáhlavský, F., Král, J., Harvey, M.S., Haddad, C.R. (2006): A karyotype study on the pseudoscorpion families Geogarypidae, Garypinidae and Olpiidae (Arachnida: Pseudoscorpiones). *European Journal of Entomology* 103:277-289.
- Tek, A.L., Jiang, J.M. (2004): The centromeric regions of potato chromosomes contain megabase-sized tandem arrays of telomere-similar sequence. *Chromosoma* 113:77-83.
- Thiry, M., Lafontaine, D.L. (2005): Birth of nucleolus: The evolution of nucleolar compartments. *Trends in Cell Biology* 15:194-199.
- Traut, W. (1976): Pachytene mapping in the female silkworm, *Bombyx mori* L. (Lepidoptera). *Chromosoma* 108:173-180.
- Traut, W., Sahara, K., Otto, T.D., Marec, F. (1999): Molecular differentiation of sex chromosomes probed by comparative genomic hybridization. *Chromosoma* 108: 173-180.
- Traut, W., Eickhoff, U., Schorch, J.C. (2001): Identification and analysis of sex chromosomes by comparative genomic hybridization (CGH). *Methods in Cell Science* 23:155-161.
- Traut, W., Szczepanowski, M., Vítková, M., Opitz, C., Marec, F., Zrzavý, J. (2007): The telomere repeat motif of basal metazoa. *Chromosome Research* 15:371-382.

- Tremousaygue, D. Manevski, A. Bardet, C. Lescure, N., Lescure, B. (1999): Plant interstitial telomere motifs participate in the control of gene expression in root meristems. *Plant Journal* 20: 553-561.
- Tres, L.L. (1975): Nucleolar RNA synthesis of meiotic prophase spermatocytes in the human testis. *Chromosoma* 53:141-151.
- Troiano G. (1990): Karyotype and male meiosis of four species of *Roncus* (Pseudoscorpionida: Neobisiidae). *Italian Journal of Zoology* 57:1-9.
- Tschochner, H., Hurt, E. (2003): Pre-ribosomes on the road from the nucleolus to the cytoplasm. *Trends in Cell Biology* 13:255-263.
- Tsuda, Y., Nishida-Umehara, Ch., Ishijima, J., Yamada, K., Matsuda, Y. (2007): Comparison of the Z and W sex chromosomal architectures in elegant crested tinamou (*Eudromia elegans*) and ostrich (*Struthio camelus*) and the process of sex chromosome differentiation in palaeognathous birds. *Chromosoma* 116:159-173.
- Tsurusaki, N. (2007): Cytogenetics. In: Pinto-da-Rocha, R., Machado, G. Giribet, G. (eds.): *Harvestmen – the biology of Opiliones*, Harvard University Press, Cambridge, USA.
- Tsurusaki, N., Cokendolpher, J.C. (1990): Chromosomes of sixteen species of harvestmen (Arachnida, Opiliones, Caddidae and Phalangidae). *Journal of Arachnology* 18:151-166.
- Tucker, S. Vitins, A., Pikaard, C.S. (2010): Nucleolar dominance and ribosomal RNA gene silencing. *Current Opinion in Cell Biology* 22:351-356.
- Uchida, W., Matsunaga, S., Sugiyama, R., Kawano, S. (2002): Interstitial telomere-like repeats in the *Arabidopsis thaliana* genome. *Genes and Genetic Systems* 77:63-67.
- Verdun, R.E., Karlseder, J. (2007): Replication and protection of telomeres. *Nature* 447:924-931.
- Vítková, M., Král, J., Traut, W., Zrzavý, J., Marec, F. (2005): The evolutionary origin of insect telomeric repeats, (TTAGG)_n. *Chromosome Research* 13:145-156.
- Vitturi, R., Colomba, M.S., Gianguzza, P., Pirrone, A.M. (2000): Chromosomal location of ribosomal DNA (rDNA), (GATA)_n and (TTAGGG)_n telomeric repeats in the neogastropod *Fasciolaria lignaria* (Mollusca: prosobranchia). *Genetica* 108: 253-257.
- Vitturi, R., Libertini, A., Armetta, F., Sparacino, L., Colomba, M.S. (2002): Chromosome analysis and FISH mapping of ribosomal DNA (rDNA), telomeric (TTAGGG)_n and (GATA)_n repeats in the leech *Haemaphysalis sanguisuga* (L.) (Annelida: Hirudinea) *Genetica* 115: 189-194.
- Wachtler, F., Stahl, A. (1993): The nucleolus: a structural and functional interpretation. *Micron* 24: 473-505.
- Walter, M.F., Bozorgnia, L., Maheshwari, A., Biessmann, H. (2001): The rate of terminal nucleotide loss from a telomere of the mosquito *Anopheles gambiae*. *Insect Molecular Biology* 10:105-110.
- Wang, Y.P., Guo, X.M. (2001): Chromosomal mapping of the vertebrate telomeric sequence (TTAGGG)_n in four bivalve molluscs by fluorescence in situ hybridization. *Journal of Shellfish Research* 20:1187-1190.
- Watson, J. D. (1972): Origin of concatemeric T7 DNA. *Nature: New Biology* 239:197-201.

Weygoldt, P., Paulus, H.F. (1979): Studies on the morphology, taxonomy and phylogeny of the Chelicerata. 2. Cladograms and development of Chelicerata. *Zeitschrift für Zoologische Systematik und Evolutionsforschung* 17:177-200.

Weygoldt, P. (2000): Whip Spiders (Chelicerata: Amblypygi): Their biology, morphology, and systematics. Apollo Books, Stenstrup, Denmark.

White, M.J.D. (1940): The origin and evolution of multiple sex chromosome mechanisms. *Journal of Genetics* 40:303-336.

White, M.J.D. (1973): *Animal cytology and evolution*, 3rd edition. Cambridge University Press, Cambridge.

Yan, J., Chen, B.Z., Bouchard, E.F., Drouin, R. (2004): The labeling efficiency of human telomeres is increased by double-strand PRINS. *Chromosoma* 113:204-209.

Yan, J., Bouchard, E.F., Samassekou, O. Chen, B.Z. (2007): Identification of a human chromosome-specific interstitial telomere-like sequence (ITS) at 22q11.2 using double-strand PRINS. *Cytogenetic and Genome Research* 116:29-37.

Zakian, V.A. (1995): Telomeres: Beginning to understand the end. *Science* 270:1601-1607.