

Errata: Study of culturable anaerobic bacterial communities living in symbiosis with bark beetles; its isolation, taxonomy and biotechnical potential.

Anna Fabryová, 2016

Chapter 4.1., page 20:

Moreover, the strains were checked for their ability to grow in aerobic conditions by placing a copy of each of the strains outside of any box.

Chapter 5.2., page 42:

Since the major molecules in plants are cellulose and hemicellulose (xylan), cellulases and xylanases are the main enzymes for plant biomass digestion, we decided to further analyse the capability of the strains of this study to synthesize cellulases and xylanases.

Chapter 8:

8. References

- Aanen, D.K., Slippers, B., Wingfield, M.J., 2009. Biological pest control in beetle agriculture. *Trends in microbiology* 17, 179–182.
- Adams, A.S., Adams, S.M., Currie, C.R., Gillette, N.E., Raffa, K.F., 2010. Geographic Variation in Bacterial Communities Associated With the Red Turpentine Beetle (Coleoptera: Curculionidae). *Environmental Entomology* 39, 406–414. doi:10.1603/EN09221
- Adams, A.S., Six, D.L., 2007. Temporal variation in mycophagy and prevalence of fungi associated with developmental stages of *Dendroctonus ponderosae* (Coleoptera: Curculionidae). *Environmental entomology* 36, 64–72.

- Adams, A.S., Six, D.L., Adams, S.M., Holben, W.E., 2008. In Vitro Interactions Between Yeasts and Bacteria and the Fungal Symbionts of the Mountain Pine Beetle (*Dendroctonus ponderosae*). *Microbial Ecology* 56, 460–466. doi:10.1007/s00248-008-9364-0
- Ayres, M.P., Wilkens, R.T., Ruel, J.J., Lombardero, M.J., Vallery, E., 2000. Nitrogen Budgets of Phloem-Feeding Bark Beetles with and without Symbiotic Fungi. *Ecology* 81, 2198. doi:10.2307/177108
- Bayer, E.A., Belaich, J.-P., Shoham, Y., Lamed, R., 2004. The Cellulosomes: Multienzyme Machines for Degradation of Plant Cell Wall Polysaccharides. *Annual Review of Microbiology* 58, 521–554. doi:10.1146/annurev.micro.57.030502.091022
- Bentz, B.J., Six, D.L., 2006. Ergosterol content of fungi associated with *Dendroctonus ponderosae* and *Dendroctonus rufipennis* (Coleoptera: Curculionidae, Scolytinae). *Annals of the Entomological Society of America* 99, 189–194.
- Berlemont, R., Martiny, A.C., 2013a. Phylogenetic Distribution of Potential Cellulases in Bacteria. *Appl. Environ. Microbiol.* 79, 1545–1554. doi:10.1128/AEM.03305-12
- Berlemont, R., Martiny, A.C., 2013b. Phylogenetic Distribution of Potential Cellulases in Bacteria. *Appl. Environ. Microbiol.* 79, 1545–1554. doi:10.1128/AEM.03305-12
- Berlemont, R., Martiny, A.C., 2015. Genomic Potential for Polysaccharide Deconstruction in Bacteria. *Appl. Environ. Microbiol.* 81, 1513–1519. doi:10.1128/AEM.03718-14
- Betts, W.B., King, J.E., 1991. Oxidative coupling of 2, 6-dimethoxyphenol by fungi and bacteria. *Mycological research* 95, 526–530.
- Brenner, D.J., Staley, J.T., Krieg, N.R., 2005. Classification of procaryotic organisms and the concept of bacterial speciation, in: *Bergey's Manual® of Systematic Bacteriology*. Springer, pp. 27–32.
- Bridges, J.R., 1981. Nitrogen-fixing bacteria associated with bark beetles. *Microb Ecol* 7, 131–137. doi:10.1007/BF02032495
- Caldwell, D.R., Bryant, M.P., 1966. Medium Without Rumen Fluid for Nonselective Enumeration and Isolation of Rumen Bacteria. *Appl Microbiol* 14, 794–801.

- Cardoza, Y.J., Moser, J.C., Klepzig, K.D., Raffa, K.F., 2008. Multipartite Symbioses Among Fungi, Mites, Nematodes, and the Spruce Beetle, *Dendroctonus rufipennis*. *Environmental Entomology* 37, 956–963. doi:10.1093/ee/37.4.956
- Cardoza, Y.J., Paskewitz, S., Raffa, K.F., others, 2006. Travelling through time and space on wings of beetles: a tripartite insect-fungi-nematode association. *SYMBIOSIS-REHOVOT*- 41, 71.
- Cardoza, Y.J., Vasanthakumar, A., Suazo, A., Raffa, K.F., 2009. Survey and phylogenetic analysis of culturable microbes in the oral secretions of three bark beetle species. *Entomologia Experimentalis et Applicata* 131, 138–147. doi:10.1111/j.1570-7458.2009.00844.x
- Case, R.J., Boucher, Y., Dahllöf, I., Holmström, C., Doolittle, W.F., Kjelleberg, S., 2007. Use of 16S rRNA and rpoB Genes as Molecular Markers for Microbial Ecology Studies. *Appl. Environ. Microbiol.* 73, 278–288. doi:10.1128/AEM.01177-06
- Cazemier, A.E., Hackstein, J.H., Den Camp, H.O., Rosenberg, J., van der Drift, C., 1997. Bacteria in the intestinal tract of different species of arthropods. *Microbial Ecology* 33, 189–197.
- Chun, J., Lee, J.-H., Jung, Y., Kim, M., Kim, S., Kim, B.K., Lim, Y.-W., 2007. EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *International journal of systematic and evolutionary microbiology* 57, 2259–2261.
- Claus, H., 2004. Laccases: structure, reactions, distribution. *Micron*, XIIIth International Conference on Invertebrate Dioxygen Binding Proteins 35, 93–96. doi:10.1016/j.micron.2003.10.029
- Collins, T., Gerday, C., Feller, G., 2005. Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiology Reviews* 29, 3–23. doi:10.1016/j.femsre.2004.06.005
- Delalibera, I., Vasanthakumar, A., Burwitz, B.J., Schloss, P.D., Klepzig, K.D., Handelsman, J., Raffa, K.F., 2007. Composition of the bacterial community in the gut of the pine engraver, *Ips pini* (Say)(Coleoptera) colonizing red pine. *Symbiosis-Rehovot*- 43, 97.
- Delorme, S., Lemanceau, P., Christen, R., Corberand, T., Meyer, J.-M., Gardan, L., 2002. *Pseudomonas lini* sp. nov., a novel species from bulk and rhizospheric soils. *International journal of systematic and evolutionary microbiology* 52, 513–523.

- Doi, R.H., Kosugi, A., 2004. Cellulosomes: plant-cell-wall-degrading enzyme complexes. *Nature Reviews Microbiology* 2, 541–551. doi:10.1038/nrmicro925
- Douglas, A.E., 2009. The microbial dimension in insect nutritional ecology. *Functional Ecology* 23, 38–47.
- Feldhaar, H., 2011. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecological Entomology* 36, 533–543.
- Furniss, M.M., Kegley, S.J., 2008. Biology of *Dendroctonus murrayanae* (Coleoptera: Curculionidae: Scolytinae) in Idaho and Montana and comparative taxonomic notes. *Annals of the Entomological Society of America* 101, 1010–1016.
- González, A.J., Tello, J.C., Rodicio, M.R., 2007. *Erwinia persicina* causing chlorosis and necrotic spots in leaves and tendrils of *Pisum sativum* in southeastern Spain. *Plant Disease* 91, 460–460.
- Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT, in: *Nucleic Acids Symposium Series*. pp. 95–98.
- Harrington, T.C., 2005. Ecology and evolution of mycophagous bark beetles and their fungal partners. *Insect-fungal associations: ecology and evolution* 1, 22.
- Hofstetter, R.W., Moser, J.C., 2014. The Role of Mites in Insect-Fungus Associations. *Annual Review of Entomology* 59, 537–557. doi:10.1146/annurev-ento-011613-162039
- Holtzapple, M., Cognata, M., Shu, Y., Hendrickson, C., 1990. Inhibition of *Trichoderma reesei* cellulase by sugars and solvents. *Biotechnol. Bioeng.* 36, 275–287. doi:10.1002/bit.260360310
- Hsiau, P.T., Harrington, T.C., 2003. Phylogenetics and adaptations of basidiomycetous fungi fed upon by bark beetles (Coleoptera: Scolytidae). *Symbiosis* 34, 111–131.
- Hu, X., Yu, J., Wang, C., Chen, H., 2014. Cellulolytic Bacteria Associated with the Gut of *Dendroctonus armandi* Larvae (Coleoptera: Curculionidae: Scolytinae). *Forests* 5, 455–465. doi:10.3390/f5030455
- Kiiskinen, L.-L., Rättö, M., Kruus, K., 2004. Screening for novel laccase-producing microbes. *Journal of applied microbiology* 97, 640–646.

- Kikuchi, Y., 2009. Endosymbiotic bacteria in insects: their diversity and culturability. *Microbes and Environments* 24, 195–204.
- Kikuchi, Y., Hosokawa, T., Fukatsu, T., 2007. Insect-microbe mutualism without vertical transmission: a stinkbug acquires a beneficial gut symbiont from the environment every generation. *Applied and environmental microbiology* 73, 4308–4316.
- Kim, O.-S., Cho, Y.-J., Lee, K., Yoon, S.-H., Kim, M., Na, H., Park, S.-C., Jeon, Y.S., Lee, J.-H., Yi, H., Won, S., Chun, J., 2012. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *International Journal Of Systematic And Evolutionary Microbiology* 62, 716–721. doi:10.1099/ijs.0.038075-0
- Kim, W.-S., Gardan, L., Rhim, S.-L., Geider, K., 1999. *Erwinia pyrifoliae* sp. nov., a novel pathogen that affects Asian pear trees (*Pyrus pyrifolia* Nakai). *International Journal of Systematic and Evolutionary Microbiology* 49, 899–906. doi:10.1099/00207713-49-2-899
- Kirisits, T., 2007. Fungal associates of European bark beetles with special emphasis on the ophiostomatoid fungi, in: *Bark and Wood Boring Insects in Living Trees in Europe, a Synthesis*. Springer, pp. 181–236.
- Krokene, P., Solheim, H., 1996. Fungal associates of five bark beetle species colonizing Norway spruce. *Can. J. For. Res.* 26, 2115–2122. doi:10.1139/x26-240
- Kube, M., Migdoll, A.M., Gehring, I., Heitmann, K., Mayer, Y., Kuhl, H., Knaust, F., Geider, K., Reinhardt, R., 2010. Genome comparison of the epiphytic bacteria *Erwinia billingiae* and *E. tasmaniensis* with the pear pathogen *E. pyrifoliae*. *BMC Genomics* 11, 393. doi:10.1186/1471-2164-11-393
- Kurtzman, C.P., 1999. Three new ascomycetous yeasts from insect-associated arboreal habitats. *Can. J. Microbiol.* 46, 50–58. doi:10.1139/w99-115
- Leufvén, A., Nehls, L., 1986. Quantification of different yeasts associated with the bark beetle, *Ips typographus*, during its attack on a spruce tree. *Microbial Ecology* 12, 237–243. doi:10.1007/BF02011208
- Lewinsohn, E., Gijzen, M., Croteau, R., 1991. Defense mechanisms of conifers differences in constitutive and wound-induced monoterpene biosynthesis among species. *Plant physiology* 96, 44–49.

- Lim, S.-H., Lee, Y.-H., Kang, H.-W., 2013. Efficient recovery of lignocellulolytic enzymes of spent mushroom compost from oyster mushrooms, *Pleurotus* spp., and potential use in dye decolorization. *Mycobiology* 41, 214–220.
- Lynd, L.R., Weimer, P.J., Zyl, W.H. van, Pretorius, I.S., 2002. Microbial Cellulose Utilization: Fundamentals and Biotechnology. *Microbiol. Mol. Biol. Rev.* 66, 506–577. doi:10.1128/MMBR.66.3.506-577.2002
- Mateos, P.F., Jimenez-Zurdo, J.I., Chen, J., Squartini, A.S., Haack, S.K., Martinez-Molina, E., Hubbell, D.H., Dazzo, F.B., 1992a. Cell-associated pectinolytic and cellulolytic enzymes in *Rhizobium leguminosarum* biovar *trifolii*. *Appl. Environ. Microbiol.* 58, 1816–1822.
- Mateos, P.F., Jimenez-Zurdo, J.I., Chen, J., Squartini, A.S., Haack, S.K., Martinez-Molina, E., Hubbell, D.H., Dazzo, F.B., 1992b. Cell-associated pectinolytic and cellulolytic enzymes in *Rhizobium leguminosarum* biovar *trifolii*. *Applied and environmental microbiology* 58, 1816–1822.
- Menéndez, E., Ramírez-Bahena, M.H., Fabryová, A., Igual, J.M., Benada, O., Mateos, P.F., Peix, A., Kolařík, M., García-Fraile, P., 2015. *Pseudomonas coleopterorum* sp. nov., a cellulase-producing bacterium isolated from the bark beetle *Hylesinus fraxini*. *International Journal of Systematic and Evolutionary Microbiology* 65, 2852–2858. doi:10.1099/ijs.0.000344
- Miao, X., Gan, M., Ding, D.-C., 2003. The role of bacterial symbionts in amino acid composition of black bean aphids. *Insect Science* 10, 167–171.
- Morales-Jiménez, J., Zúñiga, G., Ramírez-Saad, H.C., Hernández-Rodríguez, C., 2012. Gut-Associated Bacteria Throughout the Life Cycle of the Bark Beetle *Dendroctonus rhizophagus* Thomas and Bright (Curculionidae: Scolytinae) and Their Cellulolytic Activities. *Microb Ecol* 64, 268–278. doi:10.1007/s00248-011-9999-0
- Moser, J.C., Konrad, H., Kirisits, T., Carta, L.K., 2005. Phoretic mites and nematode associates of *Scolytus multistriatus* and *Scolytus pygmaeus* (Coleoptera: Scolytidae) in Austria. *Agricultural and Forest Entomology* 7, 169–177. doi:10.1111/j.1461-9555.2005.00261.x
- Nallathambi Gunaseelan, V., 1997. Anaerobic digestion of biomass for methane production: A review. *Biomass and Bioenergy* 13, 83–114. doi:10.1016/S0961-9534(97)00020-2

- Nardon, P., Grenier, A.M., 1989. Endocytobiosis in Coleoptera: biological, biochemical, and genetic aspects. *Insect endocytobiosis: morphology, physiology, genetics, evolution*. Schwemmler, W. & Gassner, G.(eds.) 175–216.
- Noda, H., Watanabe, K., Kawai, S., Yukuhiro, F., Miyoshi, T., Tomizawa, M., Koizumi, Y., Nikoh, N., Fukatsu, T., 2012. Bacteriome-associated endosymbionts of the green rice leafhopper *Nephotettix cincticeps* (Hemiptera: Cicadellidae). *Applied entomology and zoology* 47, 217–225.
- Ouhdouch, Y., Barakate, M., Finance, C., 2001. Actinomycetes of Moroccan habitats: Isolation and screening for antifungal activities. *European Journal of Soil Biology* 37, 69–74. doi:10.1016/S1164-5563(01)01069-X
- Paine, T.D., Hanlon, C.C., 1994. Influence of oleoresin constituents from *Pinus ponderosa* and *Pinus jeffreyi* on growth of mycelial fungi from *Dendroctonus ponderosae* and *Dendroctonus jeffreyi*. *J Chem Ecol* 20, 2551–2563. doi:10.1007/BF02036191
- Paine, T.D., Raffa, K.F., Harrington, T.C., 1997. Interactions among scolytid bark beetles, their associated fungi, and live host conifers. *Annual review of entomology* 42, 179–206.
- Pointing, S.B., 1999. Qualitative methods for the determination of lignocellulolytic enzyme production by tropical fungi. *Fungal Diversity*.
- Popa, V., Déziel, E., Lavallée, R., Bauce, E., Guertin, C., 2012. The complex symbiotic relationships of bark beetles with microorganisms: a potential practical approach for biological control in forestry. *Pest. Manag. Sci.* 68, 963–975. doi:10.1002/ps.3307
- Raffa, K.F., Phillips, T.W., Salom, S.M., 1993. Strategies and mechanisms of host colonization by bark beetles. *Beetle-pathogen interactions in conifer forests* 102–28.
- Richmond, C., Ujor, V., Ezeji, T.C., 2012. Impact of syringaldehyde on the growth of *Clostridium beijerinckii* NCIMB 8052 and butanol production. *3 Biotech* 2, 159–167.
- Roberts, P., 1974. *Erwinia rhapontici* (Millard) Burkholder associated with pink grain of wheat. *Journal of Applied Bacteriology* 37, 353–358.

- Ruiz-Dueñas, F.J., Martínez, Á.T., 2009. Microbial degradation of lignin: how a bulky recalcitrant polymer is efficiently recycled in nature and how we can take advantage of this. *Microbial Biotechnology* 2, 164–177. doi:10.1111/j.1751-7915.2008.00078.x
- Saini, S., Verma, A., Kumar, A., Prakash, A., Sharma, S.K., Ramesh, A., Johri, B.N., n.d. Identification and Characterization of antifungal metabolite producing *Pseudomonas protegens* strain BNJ-SS-45 isolated from rhizosphere of wheat crop (*Triticum aestivum* L.).
- Shallom, D., Shoham, Y., 2003. Microbial hemicellulases. *Current Opinion in Microbiology* 6, 219–228. doi:10.1016/S1369-5274(03)00056-0
- Shrestha, R., Koo, J.-H., Park, D.-H., Hwang, I.-G., Hur, J.-H., Lim, C.-K., 2003. *Erwinia pyrifoliae*, a Causal Endemic Pathogen of Shoot Blight of Asian Pear Tree in Korea. *The Plant Pathology Journal* 19, 294–300. doi:10.5423/PPJ.2003.19.6.294
- Sivaramakrishnan, S., Gangadharan, D., Nampoothiri, K.M., Soccol, C.R., Pandey, A., 2006. α -Amylases from microbial sources—an overview on recent developments. *Food Technol Biotechnol* 44, 173–184.
- Skrodenyte-Arbaciauskiene, V., Radziute, S., Stunzenas, V., Buda, V., 2012. *Erwinia typographi* sp. nov., isolated from bark beetle (*Ips typographus*) gut. *International Journal Of Systematic And Evolutionary Microbiology* 62, 942–948. doi:10.1099/ijss.0.030304-0
- Stark, R.W., 1982. Generalized ecology and life cycles of bark beetles. Pages 21-45 in JB Mitton & KB Sturgeon (eds) *Bark Beetles in North American Conifers: A System for the Study of Evolutionary Biology*, University of Texas Press, Austin.
- Sunna, A., Antranikian, G., 1997. Xylanolytic Enzymes from Fungi and Bacteria. *Critical Reviews in Biotechnology* 17, 39–67. doi:10.3109/07388559709146606
- Tholen, A., Schink, B., Brune, A., 1997. The gut microflora of *Reticulitermes flavipes*, its relation to oxygen, and evidence for oxygen-dependent acetogenesis by the most abundant *Enterococcus* sp.1. *FEMS Microbiology Ecology* 24, 137–149. doi:10.1016/S0168-6496(97)00053-6
- Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.* 22, 4673–4680. doi:10.1093/nar/22.22.4673

- Velicer, G.J., 2003. Social strife in the microbial world. *Trends in Microbiology* 11, 330–337. doi:10.1016/S0966-842X(03)00152-5
- Viikari, L., 1991. Xylanase enzymes promote pulp bleaching. *Paperi ja puu* 73, 384–389.
- Warren, R.A.J., 1996. Microbial Hydrolysis of Polysaccharides. *Annual Review of Microbiology* 50, 183–212. doi:10.1146/annurev.micro.50.1.183
- Wayne, L.G., Brenner, D.J., Colwell, R.R., Grimont, P.A.D., Kandler, O., Krichevsky, M.I., Moore, L.H., Moore, W.E.C., Murray, Rge., Stackebrandt, E., 1987. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *International Journal of Systematic and Evolutionary Microbiology* 37, 463–464.
- Yang, J.L., Lou, G., Eriksson, K.-E., 1992. The impact of xylanase on bleaching of kraft pulps. *Tappi journal* 75, 95–101.