

Univerzita Karlova
Přírodovědecká fakulta

Studijní program: Klinická a toxikologická analýza



Bc. Polina Yershova

**Studium redoxních a adsorpčních vlastností žlučových kyselin
na rtuťové visící kapkové elektrodě**

Study of redox and adsorption features of bile acids on hanging mercury
drop electrode

Diplomová práce

Vedoucí práce: doc. RNDr. Karolina Schwarzová, Ph.D.

Konzultant: Mgr. Lukáš Fojt, Ph.D.

Praha, 2020

Opravný list

Abstract

Bile acids are the end products of cholesterol metabolism and are important biological surfactants. The curved shape of their chains allows the cyclization of molecules, and the formation of a supramolecular structure.

The goal of this thesis was to study the electrochemical and adsorption behavior of selected bile acids: lithocholic, deoxycholic and cholic acids. The measurements were carried out in the medium Britton-Robinson buffer:methanol in the ratio 9:1 using cyclic voltammetry and AC voltammetry methods and measuring the dependence of the differential capacitance C_d on the applied potential E . A hanging mercury drop electrode was used as a working electrode.

The measurements showed that bile acids are adsorbed on the surface of the electrode and organizing themselves in self assembled monolayers (SAM). In our case we have observed formation of 2D condensed layers as specific form of SAM. Transfer techniques were used to demonstrate bile acid adsorption.

A study of the behavior of lithocholic acid as a function of different pH values showed that only at pH 10.0 to 12.0 2D condensation occurs, i. e. that at pH values in the range of 2.0 to 9.0 it is another type of adsorption. On AC voltammograms, there are a maximum of two areas in which peaks occur: the first is around -0.2 V and the second is around -1.2 V to -1.4 V. Both peaks are comparable to the peaks on a cyclic voltammogram.

The measurement of the temperature dependence of the differential capacity on the potential was performed for three concentrations of bile acids (c (mol l^{-1}) = $5 \cdot 10^{-5}$, $1 \cdot 10^{-4}$, $5 \cdot 10^{-4}$). 2D condensation in LCA occurs for both higher concentrations. At a concentration of $5 \cdot 10^{-5}$ mol l^{-1} , 2D condensation is only indicative at the lowest temperatures, but strong adsorption is still observable. DCA and CA are adsorbed on the electrode, but 2D condensation occurs only for lower temperatures at higher concentrations.

A study of the effect of the presence of anions in the basic electrolyte on the adsorption behavior of bile acids showed that the interaction of bile acids with the electrode surface and with each other is different from hydrogen bonds. Due to the structure of bile acids, it can be assumed that these are hydrophobic interactions.

All measurements showed that 2D condensation in LCA is more strong than in DCA and CA.

Cit. 10 Carey M.C., Danielsson H., Sjovall J. v knize: *Sterols and Bile Acids*. Elsevier, Amsterdam, 231 – 233 (1985).

Cit. 25 Kozoni V.; Tsioulis G.; Shiff S.; Rigas B.. *The effect of lithocholic acid on proliferation and apoptosis during the early stages of colon carcinogenesis: Differential effect on apoptosis in the presence of a colon carcinogen. Carcinogenesis* 21(5), s. 999–1005 (2000).

Cit. 26 Goldberg A. A.; Beach A.; Davies G. F.; Harkness T. A.; Leblanc A.; Titorenko V. I.. *Lithocholic bile acid selectively kills neuroblastoma cells, while sparing normal neuronal cells. Oncotarget* 2(10), s. 761–782 (2011).

Cit. 27 Ishizawa M.; Matsunawa M.; Adachi R.; Uno S.; Ikeda K.; Masuno H.; Shimizu M.: *Lithocholic acid derivatives act as selective vitamin D receptor modulators without inducing hypercalcemia. The Journal of Lipid Research* 49(4), s. 763–772 (2008).

Cit. 49 Araki Y., Mukaisyo K., Sugihara H., Fujiyama Y., Hattori T.: *Detection of N-nitroso-bile acids at 285 nm in reverse-phase HPLC. Journal of Separation Science* 31 (15), 2827–2830 (2008).

Cit. 54 Klouda J., Nesmerak K., Kocovsky P., Barek J., Schwarzova-Peckova K.: *A novel voltammetric approach to the detection of primary bile acids in serum samples, Bioelectrochemistry*, 134, 107539 (2020).

Cit. 71 Fojt L., Vetterl V., Doneux T.: *Two-dimensional condensation of nucleobases: a comparative study of halogen derivatives of cytosine. Collections of Czechoslovak Chemical Communications*, 74, 1611-1622 (2009).

Cit. 76 Fojt L., Doneux Th., Vetterl V.: *Adsorption and two-dimensional condensation of 5-methylcytosine. Bioelectrochemistry*, 75, 89-94 (2009).