

Title: Synthesis and catalytic properties of ferrocenophane phosphines

Author: Karel Škoch

Institution: Faculty of Science, Charles University in Prague, Department of Inorganic Chemistry

Supervisor: prof. RNDr. Petr Štěpnička, Ph.D.

Keywords: ferrocene, ferrocenophane, phosphine ligands, palladium, asymmetric catalysis, aza-Morita-Baylis-Hillman reaction, asymmetric allylic alkylation

Abstract: This Thesis describes the preparation of five sterically and electronically different ferrocene phosphines, (*R*)-1,1'-[1-(diarylphosphino)propan-1,3-diyl]ferrocenes (*R*)-**1a-e**, and a study into their coordination and catalytic properties. The key precursor of the phosphine synthesis, chiral alcohol (*R*)-**2**, was prepared according to the procedure described in the literature. Alcohol (*R*)-**2** was converted with retention of configuration to diarylphosphines (*R*)-**1a-e** in one-step reaction with trimethylsilylchloride and sodium iodide and then with the corresponding diarylphosphine. Phosphines **1a-e** were characterized by NMR and MS methods. For the basic representative **1a** the following palladium(II) complexes were prepared: $[\text{PdCl}(\text{L}^{\text{NC}})(\mathbf{1a})]$ (**10**, $\text{L}^{\text{NC}} = 2-[(\text{dimethylamino})\text{methyl}]\text{phenyl-}C^1,N$) and *trans*- $[\text{PdCl}_2(\mathbf{1a})_2]$ (**9a**). In addition, the isomeric complex *cis*- $[\text{PdCl}_2(\mathbf{1a})_2]$ (**9b**) was isolated from the reaction mixture after catalytic tests in asymmetric allylic alkylation. Monocrystals suitable for X-ray diffraction analysis were grown for all these complexes and the results of structure determination are presented. Phosphines (*R*)-**1a-e** were tested in two mechanistically different reactions. As organocatalysts they were used in aza-Morita-Baylis-Hillman reaction. The best results were achieved with phosphine **1a** (enantioselectivity up to 97% but low chemoselectivity). In Pd-catalyzed asymmetric allylic alkylation, the electron rich phosphine **1e** performed best achieving 98% yield and 50 % ee.