

ABSTRACT WORK

The aim of this work is to explore and describe the problem of nicotine clients of residential facilities voluntarily hospitalized for alcohol dependence. Learn whether it is on their side for such treatment if they are interested or might acquire in the course of his treatment of alcoholism motivation to quit smoking if they were offered such treatment if they were on a method of treating at least generally aware and if you know the benefits of such léčby. První stage research will consist of obtaining the basic sample. To obtain the respondents will choose the method of choice through deliberate purpose institutions. The population intended for actual research facility which clients will be allowed to participate in such research and who will be able to participate in research. Through the simple purpose a selection, Fagerström test for all survey respondents. Based on the results of this test determine how many people hospitalized for alcohol dependence with the diagnosis F10.2 simultaneously diagnoses F17.2-behavioral disorders due to use of tobacco. This finding generates the sample with which I will continue to work. To work with a sample of clients will be the main data collection method used questionnaire method. The questionnaire will be represented in both open and closed questions that the respondent offers several options. To illustrate this research will use several short case studies drawn up on the basis of interviews with specific clients using literal interpretation. In the proposed work according to the selected research methods and approaches (a deliberate choice of a special-purpose institutions through simple purposeful selection) created limits to the nature of the selected methods. Nevertheless, this work is beneficial because it opens up somewhat neglected topic of the two concurrent addictions and points to the need address this issue more than marginal attention